

Effect of some root associative bacteria on germination of seeds, nitrogenase activity and dry matter production by rice plants

A. C. DAS AND S. C. KOLE

Department of Agricultural Chemistry and Soil Science, B.C.K.V., Mohanpur, W.B.

ABSTRACT

An experiment was conducted under laboratory conditions with five efficient strains of root associative bacteria of which two N₂-fixing bacteria viz. *Azotobacter* sp. (strain AS₂) and *Azospirillum* sp. (strain AM₃), one phosphate solubilizing *Bacillus* sp. (strain BP₃), one sulphur oxidizing *Thiobacillus* sp. (strain BT₁) and one sulphur mineralizing *Bacillus* sp. (BC₅) isolated from the root surface of rice (*Oryza sativa* L. cv. IR-36) to investigate the effect of combined inoculation of the non-diazotrophic root associative bacteria (BP₃, BT₁ and BC₅) in presence of N₂-fixing bacteria on the rate of germination of rice seeds, nitrogenase activity (C₂H₂ reduction) of the roots and dry matter accumulation of the rice seedlings grown in a synthetic medium for 30 days. Inoculation of non-diazotrophic root associative bacteria in combination with N₂-fixing bacteria significantly stimulated the rate of germination of seeds as well as the nitrogenase activity of the microbes present in the roots and dry matter accumulation of rice seedlings. After 10 days of inoculation, the maximum seed germination (96.3%) was recorded with the combined inoculation of AS₂, AM₃, BP₃ and BT₁. This was followed by the combined inoculation of AS₂, BP₃, BT₁ (92.7%); AS₂, BP₃, BT₁, BC₅ (92.3%); AS₂, BT₁ (90.7%); AM₃, BP₃, BT₁ and BC₅ (90.3%), respectively. Nitrogenase activity of roots and dry matter production of the 30 day old seedlings were increased concurrently and the effects were more pronounced with the combined inoculations of AS₂, AM₃, BP₃, BT₁ followed by AS₂, BP₃, BT₁; AS₂, BP₃, BT₁, BC₅ and AM₃, BP₃, BT₁, BC₅.

Key words: Diazotrophs, Dry matter production, Germination of seeds, Inoculation effect, Nitrogenase activity, Phosphate solubilizer, Rice, Root associative bacteria, Sulphur bacteria, Synthetic medium.

The root system of any higher plant is closely associated with a vast community of active microorganisms which interact with each other for their metabolites. Consequently the fertility of soil and production of the crops are significantly influenced by the proliferation and metabolism of these active microorganisms present in the rhizosphere (Alexander, 1978). In rice rhizosphere, the availability of major plant nutrients viz. nitrogen, phosphorus and sulphur, is greatly monitored by the proliferation and activities of N₂-fixing bacteria (Das and Saha, 2005), phosphate solubilizing microorganisms (Das *et al.*, 2003) and sulphur oxidizing/mineralizing microorganisms (Saha *et al.*, 1995) closely associated with the rice roots. Besides transformation of plant nutrients, the rhizosphere microorganisms also elaborate different growth promoting substances (Arshad and Frankenberger, 1998) which are preferably utilized by the associated microbes as well as plant roots for their growth and metabolism. In the present study, an experiment has been

conducted to investigate the inoculation effect of two non-symbiotic N₂-fixing bacteria (*Azotobacter* and *Azospirillum*) either alone or in combination with phosphate solubilizing, sulphur oxidizing and sulphur mineralizing bacteria on germination of seeds, nitrogenase activity (C₂H₂ reduction) of the microbes present in the roots and dry matter accumulation by the rice plants grown on a synthetic medium under laboratory conditions.

MATERIALS AND METHODS

Five efficient strains of root associative bacteria viz. *Azotobacter* sp. (strain AS₂), *Azospirillum* sp. (strain AM₃) as non-symbiotic N₂-fixing bacteria, *Bacillus* sp. (strain BP₃) as a phosphate solubilizing bacterium, *Thiobacillus* sp. (strain BT₁) and *Bacillus* sp. (BC₅) as sulphur oxidizing and mineralizing bacteria respectively, were isolated from the root surface of rice (*Oryza sativa* L. cv. IR-36) following the methods as outlined by Das and Saha (2003). The bacterial strains were purified in their respective media through

repeated sub-culturing and microscopic examination and were identified up to their generic level following the guide of Skerman (1967) as modified by Bowie *et al.* (1969).

Rice seeds (*Oryza sativa* L. cv. IR-36) previously surface sterilized with 0.1% mercuric chloride followed by alternate washing with sterile distilled water and 70% ethyl alcohol, were inoculated with the diazotrophs (strains AS₂ and AM₃) either alone or in combination with the other non-diazotrophic root associative bacteria (BP₃, BT₁ and BC₅) by dipping the seeds in heavy suspensions (containing more than 10⁹ cells/ml) of the bacterial cultures for 1 hour followed by drying in shade. The inoculated seeds were then placed on sterile wet filter paper in sterile petriplates and were incubated at 25 ± 1°C temperature and 95 ± 1% humidity in presence of sufficient light in a growth chamber (Bittencourt *et al.*, 1995) for 10 days. There were all together 27 treatments including the uninoculated control. Each treatment was replicated thrice. The number of germinated seeds was counted after 3, 5, 7 and 10 days of incubation.

An agar tube culture experiment with the germinated rice seeds was conducted with the inoculation of efficient bacterial strains, having the same treatment as stated above. After germination, one seedling from each treatment was inoculated with the respective inoculants combinations by dipping rice roots in bacterial suspensions as stated above for 1 hour followed by air drying in shade for 30 minutes. Seedlings inoculated with bacterial cultures were then transplanted separately in 50 ml sterile semisolid plant nutrient agar medium (Murasighe and Skoog, 1962) in culture tubes under laboratory conditions. The composition of the medium was (per liter): NH₄NO₃, 1.65 g; KNO₃, 1.9 g; CaCl₂·2H₂O, 440 mg; MgSO₄·7H₂O, 370 mg; KH₂PO₄, 170 mg; KI, 0.83 mg; H₃BO₄, 6.2 mg; MnSO₄·4H₂O, 22.3 mg; ZnSO₄·7H₂O, 8.6 mg; Na₂MoO₄·2H₂O, 0.25 mg; CaCl₂·6H₂O, 0.025 mg; FeSO₄·7H₂O, 27.8 mg; CuSO₄·5H₂O, 0.025 mg; Na₂-EDTA, 37.3 mg; inositol, 100 mg; nicotinic acid, 0.5 mg; pyridoxine-HCl, 0.5 mg; thiamin, 0.1 mg; IAA, 1-30 mg; kinetin, 0.4 -10 mg; glycine, 2 mg; sucrose, 30

g and agar 8 g. The pH of the medium was adjusted to 5.7. After 30 days of transplanting, the seedlings were carefully uprooted from the medium and the roots were washed with sterile distilled water. Thereafter, nitrogenase activity of the roots was determined through acetylene reduction assay (ARA) (Ghosh and Saha, 1993), with the help of a gas chromatograph (HP model 5730A) fitted with a glass column packed with porapak R (80-100 mesh) and equipped with a flame ionizing detector. The operating temperature of the oven and the flow rate of carrier gas (N₂) were adjusted to 80°C and 60 ml/min, respectively. The dry matter content of the 30-day old rice plants was also determined following standard methods.

RESULTS AND DISCUSSION

Effect on seed germination

Inoculation of *Azotobacter* (strain AS₂) and *Azospirillum* (strain AM₃) singly or in combination with the phosphate solubilizing *Bacillus* (strain BP₃), sulphur oxidizing *Thiobacillus* (strain BT₁) and sulphur mineralizing *Bacillus* (strain BC₅), significantly stimulated the germination of rice seeds as compared to uninoculated control (Table 1). The bacterial combinations AS₂, AM₃, BP₃, BT₁ and BC₅ significantly increased the germination of seeds from 3rd day onwards. In general, the maximum germination of seeds was recorded with the inoculation of the bacterial strains after 5 and 7 days of inoculation and the combined inoculation of phosphate solubilizing *Bacillus* (strain BP₃), sulphur oxidizing *Thiobacillus* (strain BT₁) and sulphur mineralizing *Bacillus* (strain BC₅) with *Azotobacter* (strain AS₂) and/or *Azospirillum* (strain AM₃) showed better effect as compared to their single inoculation. This indicated that non-symbiotic N₂-fixing bacteria along with other root associative non-diazotrophic bacteria (strains BP₃, BT₁ and BC₅) increased the availability of nitrogen, phosphorus and sulphur as well as some growth promoting substances released by the microorganisms (Park *et al.*, 2005) in the vicinity of the seeds which subsequently stimulated the germination of rice seeds to a greater extent. It was also revealed that the effect of the diazotrophs (strains AS₂ and AM₃) on seed germination

was more pronounced when they were in association with the other root associative non-diazotrophic bacteria responsible for releasing mineral phosphorus, sulphur and other growth promoting substances for the growth and metabolism of the rhizosphere microorganisms as well as the young seedlings germinating from the inoculated seeds. Similar observations were also reported by earlier workers (Thakuria *et al.*, 2004). After 10 days of inoculation, the maximum seed germination (96.3%) was recorded with the combined inoculation of AS₂ + AM₃ + BP₃ + BT₁. This was followed by the combined inoculation of AS₂ + BP₃ + BT₁ (92.7%), AS₂ + BP₃ + BT₁ + BC₅ (92.3%), AS₂ + BT₁ (90.7%) and AM₃ + BP₃ + BT₁ + BC₅ (90.3%), respectively.

Effect on nitrogenase activity (C₂H₂ reduction)

Inoculation of *Azotobacter* (strain AS₂) and *Azospirillum* (strain AM₃) either alone or in combination with phosphate solubilizing *Bacillus* (strain BP₃), sulphur oxidizing *Thiobacillus* (strain BT₁) and sulphur mineralizing *Bacillus* (strain BC₅) increased the nitrogenase activity (C₂H₂ reduction) of the microbes present in the roots of the rice seedlings as compared to the uninoculated control (Table 2) and the stimulation was more pronounced when the diazotrophs were inoculated in combination with the other non-diazotrophic root associative bacteria (BP₃, BT₁ and BC₅) rather than their single inoculation. This indicated that phosphate solubilizing and sulphur oxidizing/mineralizing bacteria had a proto-cooperative association with the non-symbiotic N₂-fixing bacteria (*Azotobacter* and *Azospirillum*) in the root rhizosphere of rice plants resulting in an increase in their activities which augmented the nitrogenase activity of the diazotrophs (strain AS₂ and AM₃) in the root zone to a great extent (Das and Saha, 2005). This was in agreement with the earlier workers (Park *et al.*, 2005) who reported that the versatile groups of microorganisms isolated from the rice rhizosphere had a significant effect on the nitrogenase activity of the rice roots when the organisms were inoculated in combination. As compared to uninoculated control, the highest increase in nitrogenase

activity was recorded (185.9%) in the rice roots when the seedlings were inoculated with AS₂, AM₃, BP₃ and BT₁ in combination. This was followed by the combined inoculation of AS₂ + BP₃ + BT₁ (165.9%), AS₂ + BP₃ + BT₁ + BC₅ (127.4%) and AM₃ + BP₃ + BT₁ + BC₅ (87.4%), respectively. It was also revealed that the effect of *Azotobacter* (strain AS₂) in association with other non-diazotrophic bacteria was more pronounced than that of *Azospirillum* (strain AM₃). This indicated that the activity of *Azotobacter* was highly stimulated due to the presence of other root associative non-diazotrophic bacteria that could have released more amounts of phosphorus, sulphur and other growth promoting substances to the diazotrophs as well as to the plants for their growth and metabolism (Park *et al.*, 2005). This supported the existence of more compatible association of the non-diazotrophic root associative bacteria with *Azotobacter* than that with *Azospirillum*.

Effect on dry matter production

Sustaining the earlier reports (Thakuria *et al.*, 2004), the dry matter production of the rice plants were concomitantly increased with the nitrogenase activity of the rice roots. This indicated that greater N₂-fixation together with greater solubilization of insoluble phosphates, oxidation of inorganic sulphur and mineralization of organic sulphur in the root rhizosphere of the rice plants by the active microbes significantly increased the availability of nitrogen, phosphorus and sulphur in the root zone (Table 2). This accelerated the metabolic activities of the plants resulting greater accumulation of dry matter content of the plant as compared to the uninoculated control during the incubation period. Among the treatments, the maximum increase in the accumulation of dry matter was recorded with the inoculation combination AS₂, AM₃, BP₃ and BT₁ (77.5%) followed by AS₂, BP₃ and BT₁ (76.1%), AS₂, BP₃, BT₁ and BC₅ (60.6%), and AM₃, BP₃, BT₁ and BC₅ (36.6%), respectively as compared to the uninoculated control.

The results of the present investigation thus clearly indicated that the inoculation of

Table 1 Effect of inoculation of diazotrophs and some root associative bacteria on the germination (in per cent) of rice seeds

Treatments	Incubation period in days			
	3	5	7	10
Control (uninoculated)	28.7	30.7	56.0	75.3
<i>Azotobacter</i> (strain AS ₂)	25.0	57.3	76.0	82.0
<i>Azospirillum</i> (strain AM ₃)	23.3	65.3	79.0	79.3
P-solubilizing <i>Bacillus</i> (strain BP ₃)	26.3	52.3	78.3	85.0
S-oxidizing <i>Thiobacillus</i> (strain BT ₁)	23.3	69.7	76.7	81.7
S-mineralizing <i>Bacillus</i> (strain BC ₅)	24.3	61.3	77.7	84.3
AS ₂ + AM ₃	29.3	73.0	79.0	82.3
AS ₂ + BP ₃	27.7	75.3	82.3	85.3
AS ₂ + BT ₁	24.0	75.7	86.7	90.7
AS ₂ + BC ₅	26.0	65.0	75.0	83.0
AM ₃ + BP ₃	22.3	67.7	73.7	84.0
AM ₃ + BT ₁	27.7	69.0	82.0	87.7
AM ₃ + BC ₅	23.3	71.0	83.0	85.7
AS ₂ + AM ₃ + BP ₃	25.0	73.7	81.7	86.7
AS ₂ + AM ₃ + BT ₁	23.3	56.7	75.3	85.0
AS ₂ + AM ₃ + BC ₅	22.7	64.0	74.6	85.7
AS ₂ + BP ₃ + BT ₁	38.0	84.0	88.7	92.7
AS ₂ + BP ₃ + BC ₅	25.7	70.7	84.3	87.7
AS ₂ + BT ₁ + BC ₅	27.0	68.0	78.0	83.0
AM ₃ + BP ₃ + BT ₁	25.0	70.0	81.0	84.7
AM ₃ + BP ₃ + BC ₅	23.0	69.0	75.0	78.3
AM ₃ + BT ₁ + BC ₅	22.0	59.3	72.7	80.0
AS ₂ + AM ₃ + BP ₃ + BT ₁	39.7	85.0	87.7	96.3
AS ₂ + AM ₃ + BP ₃ + BC ₅	25.0	63.7	73.0	79.7
AS ₂ + BP ₃ + BT ₁ + BC ₅	29.3	77.3	85.7	92.3
AM ₃ + BP ₃ + BT ₁ + BC ₅	26.0	64.3	81.0	90.3
AS ₂ + AM ₃ + BP ₃ + BT ₁ + BC ₅	22.7	65.0	78.0	85.7
Mean	26.1	66.8	78.6	84.9
CD at 5%	Treatments 3.3; Incubation days 1.9; Interaction 10.1			

Table 2 Effect of inoculation of diazotrophs and some root associative bacteria on nitrogenase activity (C₂H₂ reduction) of roots and dry matter production of rice seedlings grown in a synthetic medium

Treatments	C ₂ H ₂ reduced by roots	Dry matter produced (30 DAS)
	(nmol/100 mg/hr)	(mg/plant)
Control (uninoculated)	135	71
<i>Azotobacter</i> (strain AS ₂)	190	99
<i>Azospirillum</i> (strain AM ₃)	173	94
P-solubilizing <i>Bacillus</i> (strain BP ₃)	162	83
S-oxidizing <i>Thiobacillus</i> (strain BT ₁)	169	87
S-mineralizing <i>Bacillus</i> (strain BC ₅)	165	88
AS ₂ + AM ₃	211	101
AS ₂ + BP ₃	144	93
AS ₂ + BT ₁	182	92
AS ₂ + BC ₅	148	77
AM ₃ + BP ₃	135	102
AM ₃ + BT ₁	183	79
AM ₃ + BC ₅	171	82
AS ₂ + AM ₃ + BP ₃	200	76
AS ₂ + AM ₃ + BT ₁	175	86
AS ₂ + AM ₃ + BC ₅	204	85
AS ₂ + BP ₃ + BT ₁	359	125
AS ₂ + BP ₃ + BC ₅	233	90
AS ₂ + BT ₁ + BC ₅	194	98
AM ₃ + BP ₃ + BT ₁	147	99
AM ₃ + BP ₃ + BC ₅	181	99
AM ₃ + BT ₁ + BC ₅	159	103
AS ₂ + AM ₃ + BP ₃ + BT ₁	386	126
AS ₂ + AM ₃ + BP ₃ + BC ₅	179	97
AS ₂ + BP ₃ + BT ₁ + BC ₅	307	114
AM ₃ + BP ₃ + BT ₁ + BC ₅	253	111
AS ₂ + AM ₃ + BP ₃ + BT ₁ + BC ₅	192	78
CD at 5%	39.3	11.2

root associative bacteria significantly stimulated the germination of seeds, nitrogenase activity (C_2H_2 reduction) of the microbes present in the roots and the dry matter accumulation by rice seedlings grown in a synthetic medium. It was also revealed that the combined inoculation of the organisms responded better than their single inoculation and the response was more pronounced when the diazotrophic bacteria (*Azotobacter* than *Azospirillum*) were inoculated in combination with the non-diazotrophic bacteria (phosphate solubilizing *Bacillus*, *Thiobacillus* and sulphur mineralizing *Bacillus*). Between the two non-symbiotic N_2 -fixing bacteria, the compatibility of the non-diazotrophic bacteria was better with *Azotobacter* than with *Azospirillum*.

REFERENCES

- Alexander M. 1978. *Introduction to Soil Microbiology*. Wiley Eastern Ltd., New Delhi, India.
- Arshad M and Frankenberger Jr W T. 1998. Plant growth-regulating substances in the rhizosphere: microbial production and function. *Advances in Agronomy*, **62**: 45-151.
- Bittencourt S R M, Vieira R D, Barreto M and Volpe C A. 1995. Comparison of two types of seed germinators for use as a chamber for the accelerated aging test. *Revista Brasileira de Sementes*, **17** (1): 69-74.
- Bowie I S, Loutit M W and Loutit J S. 1969. Identification of aerobic heterotrophic soil bacteria to the generic level by using multipoint inoculation techniques. *Canadian Journal of Microbiology*, **15**: 297-302.
- Das A C and Saha D. 2003. Influence of diazotrophic inoculations on nitrogen nutrition of rice. *Australian Journal of Soil Research*, **41** (8): 1543-1554
- Das A C and Saha D. 2005. Non-symbiotic nitrogen-fixing bacteria influencing mineral and hydrolysable nitrogen in rhizosphere soils of rice (*Oryza sativa*). *Indian Journal of Agricultural Sciences*, **75** (5): 265-269.
- Das A C, Debnath A and Mukherjee D. 2003. Effect of the herbicides oxadiazon and oxyfluorfen on phosphates solubilizing microorganisms and their persistence in rice fields. *Chemosphere*, **53** (3): 217-221.
- Ghosh T K and Saha K C. 1993. Effect of inoculation with N_2 -fixing cyanobacteria on the nitrogenase activity in soil and rhizosphere of wetland rice (*Oryza sativa* L). *Biology and Fertility of Soils*, **16** (1): 16-20.
- Murasighe T and Skoog F. 1962. A revised medium for rapid growth and bioassay with tobacco tissue culture. *Physiologia Plantarum*, **15**: 473-497.
- Park M S, Kim C W, Yang J C., Lee H S, Shin W S, Kim S H and Sa T M. 2005. Isolation and characterization of diazotrophic growth promoting bacteria from rhizosphere of agricultural crops of Korea. *Microbiological Research*, **160** (2): 127-133.
- Saha N, Das A C and Mukherjee D. 1995. Effect of decomposition of organic matter on the activities of microorganisms and availability of nitrogen, phosphorus and sulphur in soil. *Journal of the Indian Society of Soil Science*, **43** (2): 210-215.
- Skerman V B D. 1967. *A Guide to the Identification of the Genera of Bacteria*. The Williams and Wilkins Company, Baltimore, USA.
- Thakuria D, Talukdar N C, Goswami C, Hazarika S, Boro R C and Khan M R. 2004. Characterization and screening of bacteria from rhizosphere of rice grown in acidic soils of Assam. *Current Science*, **86** (7): 978-985.