

Effect of sulphur on growth and yield of green gram [*Vigna radiata* (L.) Wilczek]

N. C. BANIK AND K. SENGUPTA

Department of Agronomy, Faculty of Agriculture
Bidhan Chandra Krishi Viswavidyalaya, Mohanpur-741252
Nadia, West Bengal, India.

Received: 10.06.2011, Revised: 17.03.2012, Accepted : 25.04.2012

ABSTRACT

A field experiment was conducted to evaluate the effect of phospho-gypsum on green gram (*Vigna radiata* L.) in Gangetic alluvial soil during two successive seasons. The experiment was laid out in RBD with six treatments and four replications. The results reveal that application of Phospho-gypsum at 30 kg S / ha with the recommended dose of N-P-K (T_3) gives highest seed yield of green gram crop followed by T_5 treatment where 30 kg S / ha automatically supplied by Single Super Phosphate with the recommended N-P-K dose and T_4 treatment where Phospho-gypsum at 20 kg S / ha applied with the recommended dose of N-P-K in both the seasons. Seed yield obtained in treatments T_5 and T_3 significantly differ from T_2 treatment where recommended fertilizer dose (N-P-K) applied without any sulphur. Crop growth characters like plant height and total dry matter production also showed similar results. Phospho-gypsum was found effective and may be used as an alternate source of S for increasing the yield of green gram crop. The dose of S 20-30 kg ha⁻¹, may be optimum and better option for increasing the yield of green gram crop in an intensive cropping system where green gram is grown as a catch crop / soil restoring crop.

Key words: Growth, net production value, phospho-gypsum, sulphur, yield

Green gram [*Vigna radiata* (L.) Wilczek], commonly known as 'Mung bean or Moong', being a legume crop has high demand for sulphur due to production of several protein containing materials and fatty acids, in which sulphur is an important constituent. Sulphur is required in plant for the synthesis of chlorophyll and essential for activation of certain proteolytic enzymes such as papainase. Increased use of sulphur-free fertilizers, intensive cropping, and use of high-yielding varieties have led to S deficiency in many countries. Sulphur deficiency is increasingly becoming one of the limiting factors to further sustainable increase in agricultural production. Sulphur fertilizer, besides enhancing yield and quality of crops, enhances nutrient uptake, particularly N, and fertilizer-use efficiency through interaction of sulphur with other fertilizer nutrients (Tandon and Messick, 2007). S is needed for conversion of reduced N into protein in symbiotic N fixation in pulses (like green gram), thus its positive effect on N absorption is quite likely. Residual effects have also been reported even at a low rate of 20 kg S ha⁻¹. Thus, sulphur does not need to be applied every season (De Datta *et al.*, 2005). Application of sulphur along with N, P and K to pulses and oilseeds showed greater response than to cereals. Sulphur not only improved grain yield but also improved the quality of crops (Hegde and Babu, 2004; Hegde and Murthy, 2005). Mitra *et al.* (2006) reported that green gram yield increased with increasing levels of P and S fertilizers. P at 60 kg P₂O₅ and S at 40 kg ha⁻¹ gave the highest number of pods per plant, number of seeds per pod, 1000-seed weight, seed yield, net return and net return/rupee invested. However, meager information is available on the effect of Phospho-gypsum as an alternate source of S for the growth and yield of green gram crop. Hence the present investigation was aimed to study the effect of S on green gram using phospho-gypsum as one of the sources.

MATERIALS AND METHODS

The experiment was undertaken during two successive seasons (autumn-2008 and summer-2009) to study the effect of sulphur on growth and yield of green gram [*Vigna radiata* (L.) Wilczek] crop (variety PDM-54). The experiment was laid out in RBD with 6 treatments and 4 replications at Instructional Farm (New Alluvial Zone) of Bidhan Chandra Krishi Viswavidyalaya, Nadia, West Bengal (situated at 22°56' N latitude, 88°32' E longitude and 9.75 m MSL. The soil characters of the experimental field were: pH 7.1, organic carbon 0.68%, total N 0.07%, available P₂O₅ 28.3 kg ha⁻¹, and available K₂O 126.5 kg ha⁻¹ and available sulphate 10.3 ppm. The treatments were: T_1 = Control (No fertilizer), T_2 = Fertilized control-I i.e. 20-40-40 N-P₂O₅-K₂O kg ha⁻¹ (Recommended fertilizer dose, without any S) P through DAP, T_3 = Fertilized control-II i.e., 20-40-40 N-P₂O₅-K₂O kg ha⁻¹ (Recommended fertilizer dose, with S) P through SSP, T_4 = T_2 + Phospho-gypsum at 20 kg S/ha, T_5 = T_2 + Phospho-gypsum at 30 kg S ha⁻¹, T_6 = T_2 + Phospho-gypsum at 10 kg S ha⁻¹. The land preparation, pre-sowing seed treatment, sowing of seeds, irrigation application and intercultural operations such as weeding, hoeing, thinning, plant protection measures and harvesting were done as per recommended practices followed in green gram and requirement of the field. Observations on the growth characters and seed yield were taken and analyses were done for the determination of treatment effect.

RESULTS AND DISCUSSION

Table-1 showed the effect of sulphur on growth and seed yield of green gram crop. Plant height and dry aerial bio-mass were recorded at 20, 40 and 60 DAS in both the seasons. The result showed that highest plant height obtained in T_5 followed by T_3 and T_4 treatment. Treatment T_5 (Recommended N-P-K dose + Phospho-gypsum at 30 kg S/ha) and T_3 (Recommended N-P-K dose, P through SSP as a result S was added

automatically at 30 kg ha⁻¹) significantly differed from T₁ (No fertilizer) and T₂ (Recommended dose of N-P-K

without any S) treatment at 20 and 40 DAS in both the seasons. Similar trend was observed at 60 DAS.

Table 1: Effect of sulphur on growth and yield of green gram crop

Treatment	Plant height (cm)						Total dry matter [dry aerial bio-mass] (kg ha ⁻¹)						Seed yield (kg ha ⁻¹)	
	Autumn-2008			Summer-2009			Autumn-2008			Summer-2009			Autumn- 2008	Summer- 2009
	20	40	60	20	40	60	20	40	60	20	40	60		
Days after sowing														
T ₁	6.9	44.6	43.6	11.1	36.3	43.5	104.1	1003.2	2197.2	61.918	1410.8	2115.5	608.5	587.6
T ₂	7.3	47.9	46.7	11.7	39.3	50.5	116.0	1068.5	2820.4	69.625	1443.5	2304.8	706.0	697.8
T ₃	8.1	52.5	54.2	12.5	43.1	52.0	131.7	1410.5	3068.9	75.915	1635.0	2666.5	1093.7	1097.5
T ₄	7.8	48.7	53.7	12.0	40.6	50.9	132.5	1344.9	2993.0	74.085	1568.3	2429.0	943.0	1072.2
T ₅	8.2	53.2	54.4	12.7	43.5	54.6	135.9	1445.7	3570.5	76.083	1705.0	2693.3	1101.3	1160.0
T ₆	6.8	48.4	51.9	11.8	39.9	49.65	129.4	1125.7	3237.2	75.003	1562.5	2306.8	915.0	1032.4
SEm (\pm)	0.22	0.95	2.11	0.23	1.25	1.99	4.11	89.84	243.01	2.07	54.07	174.44	89.0	104.4
LSD(0.05)	0.65	2.87	6.36	0.71	3.76	6.01	12.38	270.81	732.51	6.25	162.98	525.83	268.4	314.7

The result indicated that highest total dry matter production obtained in T₅ treatment followed by T₃, T₄ and T₆ treatments (Table 1). Total dry matter production in T₅ and T₃ treatments significantly differed from T₂ treatment at 20 and 40 DAS in both the seasons. At 60 DAS total dry matter production in T₅ treatment was significantly better than T₂ treatment in both the seasons. Umesh *et al.* (1997) reported similar results that application of S at 30 kg ha⁻¹ was optimum for plant height, branch number, functional leaves, dry matter accumulation, dry weight of nodules/plant, and seed yield.

Table- 1 showed seed yield of green gram crop varied due to different level of sulphur application. The results revealed that application of Phospho-gypsum at 30 kg S ha⁻¹ with the recommended dose of N-P-K (T₅) gave highest seed yield of green gram crop followed by T₃ treatment where S at 30 kg ha⁻¹ was added automatically through Single Super Phosphate with the recommended N-P-K dose and T₄ treatment where Phospho-gypsum at 20 kg S / ha applied with the recommended dose of N-P-K in both the seasons. Seed yield obtained in treatments T₅ and T₃ significantly differed from T₂ treatment where recommended fertilizer dose (N-P-K) applied without any sulphur. Jat and Rathore (1994) reported that seed yield of green gram increased with sulphur application up to 30 kg ha⁻¹. Singh and Kumar (1996) also reported that application of S at 30 kg ha⁻¹ significantly increased the pods per plant, 1000 seed weight and also gave highest seed and straw yields of lentil.

The net production values (NPV) of green gram crop raised in two seasons were studied. Application of nutrients from external source gave remarkably higher net profit over absolute control treatment (T₁) where no fertilizer was applied. Maximum net production values of 2.60 recorded in the treatment T₃, receiving sulphur from SSP (used as P source), followed by treatment T₅ (NPV 2.29), receiving phospho-gypsum at 30 kg S ha⁻¹, and T₄ (NPV 1.91), receiving phospho-gypsum at 20 kg S ha⁻¹. Application of S through any source increased the NPV over fertilized control treatment.

Sulphur was found to be effective on growth and yield of green gram crop. Result showed that growth and yield of green gram crop increased significantly with N-P-K applied along with sulphur through Phospho-gypsum than the crop receiving no sulphur. Therefore, Phospho-gypsum may be used as an alternate source of S for increasing the yield of green gram crop. The dose of S at 20 or 30 kg ha⁻¹ may be optimum and better option for increasing the yield of green gram crop in an intensive cropping system where green gram is grown as a catch crop or soil restoring crop.

REFERENCES

- De Datta, S. K., Buresh, R. J. and Mamaril, C. P. 2005. Increasing nutrient-use efficiency in rice with changing needs. *Nutrient Cycling in Agro-ecosystems*, 26: 157-67.
- Hegde, D. M. and Murthy, I.Y.L.N. 2005. Management of secondary nutrients: achievements and challenges. *Indian J. Fertilizers*, 1: 93-10.
- Hegde, D.M. and Sudhakara Babu, S.N. 2004. Role of balanced fertilization in improving crop yield and quality. *Fertilizer News*, 49: 103-31
- Jat, R.L. and Rathore, P.S. 1994. Effect of sulphur, molybdenum and Rhizobium inoculation on greengram (*Phaseolus radiatus*). *Indian J. Agron.*, 39: 651-54.
- Mitra, A.K., Banerjee, K. and Pal, A. K. 2006. Effect of different levels of phosphorus and sulphur on yield attributes, seed yield, protein content of seed and economics of summer greengram. *Res. Crops*, 7: 404-05.
- Singh, B. and Kumar, V. 1996. Effect of P and S on lentil (*Lens culinaris*) under rainfed condition. *J. Agron.*, 41: 420-23.
- Tandon, H. L. S. and Messick, D. L. 2007. *Practical Sulphur Guide*. The Sulphur Institute, Washington, USA.
- Umesh, Singh., Yadav, D.S. and Singh, U. 1997. Studies on sulphur and zinc nutrition on greengram (*Phaseolus radiatus* L.) in relation to growth attributes, seed protein yield and S, Zn uptake. *Legume Res.*, 20: 224-26.