

## Effect of different planting patterns and fertilizer levels on production potential of maize [*Zea mays* L.] and green gram (*Vigna radiata* L.)

P. B. BEHERE, V. H. SURVE, R.R. PISAL, <sup>1</sup>P.R. PATIL AND V.C. RAJ

Department of Agronomy and <sup>1</sup>Department of Plant Breeding  
N. M. College of Agriculture, Navsari Agricultural University,  
Navsari-396450, Gujarat,

Received:04-08-2013, Revised:20-11- 2013, Accepted: 22-11-2013

**Key words:** Intercropping, economics, fodder yield and planting pattern

Sweet corn (*Zea mays* L) is one of the most important cereals, which ranks fifth in India for production. Maize is grown in 6.5 million ha in India and 45 per cent of the total production is consumed as food. It is used as food, feed and forage as well as in industry. India is the homeland of grain legumes including pulses which play vital role in cereal based Indian diet. Green gram (*Vigna radiata* L.) locally known as 'Moong' in India covering an area of 3.1 million hectares with a total production of 1.1 million tonnes with the average of 425 ha<sup>-1</sup> (Anon., 2004). Intercropping has been recognised as a beneficial system of crop production and is one of the potent means of better utilization of resources and higher crop production per unit time and area, which can provide substantial yield advantages compared to sole cropping. These advantages may be specially important because they are achieved not by means of costly inputs, but by the simple expedient of growing crop together (Willey, 1979). In new system, the modification of planting geometry may helps in accommodating the companion crop. By adopting the appropriate planting pattern the total productivity can be enhanced.

The field experiment was conducted during summer season of 2007 at the College of Agriculture, Navsari, Gujarat. The soil of the experimental field was clayey in texture, low in nitrogen (201 ha<sup>-1</sup>), medium in available phosphorus (30.52 ha<sup>-1</sup>) and fairly rich in available potassium (352 ha<sup>-1</sup>) with pH (7.8). Nine treatment combinations consisting of three planting pattern and various nitrogen levels T<sub>1</sub>: Sole sweet corn uniform row of 60 cm (control), T<sub>2</sub>: Sole green gram uniform row of 30 cm (control), T<sub>3</sub>: Paired row normal planting (control), T<sub>4</sub>: Sweet corn + green gram (1:1) uniform in row of 60 cm (75 % RFD of N in sweet corn), T<sub>5</sub>: Sweet corn + green gram (1:1) uniform in row of 60 cm (100 % RFD of N in sweet corn), T<sub>6</sub>: Sweet corn + green gram (1:1) uniform in row of 60 cm (125 % RFD of N in sweet corn), T<sub>7</sub>: Sweet corn + green gram (2:2) in paired row of 45/90 cm (75 % RFD of N in sweet corn), T<sub>8</sub>: Sweet corn + green gram (2:2) in paired row of 45/90 cm (100 % RFD of N in sweet corn) T<sub>9</sub>: Sweet corn + green gram (2:2) in paired row of 45/90 cm (125 %

RFD of N in sweet corn) were evaluated in randomized block design with four replications. The varieties *Madhuri* and GM-4 respectively for sweet corn and green gram were sown on last week of Feb, 2007. The seed rate under sole cropping was maintained at 20 and 12 ha<sup>-1</sup> respectively for maize and green gram.

**Table 1: Cob/ grain and fodder yields and harvest index in sweet corn and green gram as influenced by planting pattern and fertility levels**

Treat ments	Yield (q.ha <sup>-1</sup> )		HI (%)	Yield (q.ha <sup>-1</sup> )		HI (%)
	Cob	Fodder		Grain	Fodder	
T <sub>1</sub>	68.48	133.56	33.89	-	-	-
T <sub>2</sub>	-	-	-	9.68	23.91	28.82
T <sub>3</sub>	51.48	103.30	33.26	3.31	8.74	27.47
T <sub>4</sub>	62.89	121.77	34.05	3.90	9.21	29.76
T <sub>5</sub>	65.15	129.74	33.43	4.38	10.01	30.44
T <sub>6</sub>	63.37	126.27	33.42	4.07	9.86	29.23
T <sub>7</sub>	62.38	126.55	33.01	3.97	9.81	28.81
T <sub>8</sub>	68.92	132.24	34.26	4.46	10.06	30.69
T <sub>9</sub>	66.37	129.34	33.91	4.08	9.89	29.19
<b>S.Em.(+)</b>	<b>0.59</b>	<b>0.74</b>	<b>0.26</b>	<b>0.11</b>	<b>0.18</b>	<b>0.65</b>
<b>LSD(0.05)</b>	<b>1.74</b>	<b>2.19</b>	<b>0.77</b>	<b>0.33</b>	<b>0.53</b>	<b>NS</b>
<b>C.V.%</b>	<b>1.87</b>	<b>1.19</b>	<b>1.56</b>	<b>4.76</b>	<b>3.21</b>	<b>4.49</b>

HI: Harvest Index

Treatments of planting pattern and fertility levels significantly influenced the yield of maize and green gram (Table 1). The treatment T<sub>8</sub> (maize + green gram (2:2) in paired row of 45/90 cm (100 % RFD of N in sweet corn) recorded significantly the highest cob yield (68.92 q ha<sup>-1</sup>) however, it was statistically at par with T<sub>1</sub>. Treatment T<sub>3</sub> (row normal planting) (control) recorded significantly the lowest cob yield (51.48 q ha<sup>-1</sup>) of sweet corn. The treatment T<sub>8</sub> and T<sub>1</sub> increased the cob yield by 33.87 and 33.02 per cent respectively, over T<sub>3</sub>. Similar results were also reported by Sharma (1995), Akhtar and Silva (1999) and Shivran and Rana (2003).

The highest fodder yield (133.56 q ha<sup>-1</sup>) was obtained under treatment T<sub>1</sub> (sweet corn uniform row of 60 cm) (control) but found at par with T<sub>8</sub>. Significantly the lowest fodder yield (103.30 q ha<sup>-1</sup>) was recorded with T<sub>3</sub> (paired row normal planting) (control). The treatment T<sub>1</sub> and T<sub>8</sub>

increased the fodder yield by 29.29 and 28.01 per cent, respectively, over T<sub>3</sub>. Similar results were also reported by Arya and Saini (1989) and Sharma (1995).

The treatment T<sub>8</sub> (corn + green gram (2:2) in paired row of 45/90 cm) (100 % RFD of N in sweet corn) recorded maximum harvest index, however, it was found *at par* with T<sub>4</sub>, T<sub>9</sub> and T<sub>1</sub>. The lowest value was observed under treatment T<sub>7</sub> (corn + green gram (2:2) in paired row of 45/90 cm) (75 % RFD of N in sweet corn) being *at par* with T<sub>3</sub>, T<sub>6</sub> and T<sub>5</sub>. Similar results were also reported by Padhi and Panigrahi (2006), The results are contradictory with findings of Dhingra *et al.* (1991).

The treatment T<sub>2</sub> (green gram uniform row of 30 cm) (control) recorded significantly the higher grain yield (9.68 q ha<sup>-1</sup>) compared to other treatments. Treatment of paired row normal planting (control) (T<sub>3</sub>), recorded significantly the lowest grain yield (3.31 q ha<sup>-1</sup>) of green gram indicating yield reduction due to normal planting pattern.

**Table 2: Economics of different intercropping systems**

Treat-ments	Gross return (₹. ha <sup>-1</sup> )	Cost of cultivation (₹. ha <sup>-1</sup> )	Net return (₹. ha <sup>-1</sup> )	BCR
T <sub>1</sub>	68140	12679	55461	5.37
T <sub>2</sub>	21751	9155	12596	2.37
T <sub>3</sub>	59008	12299	46709	4.79
T <sub>4</sub>	71210	13693	57517	5.20
T <sub>5</sub>	74855	14126	60729	5.29
T <sub>6</sub>	72449	14459	57990	5.01
T <sub>7</sub>	71480	13558	57922	5.27
T <sub>8</sub>	78286	13891	64395	5.63
T <sub>9</sub>	75171	14224	60947	5.28

Treatment receiving T<sub>2</sub> (green gram uniform row of 30 cm) (control) produced the highest fodder yield (23.91 q ha<sup>-1</sup>). Treatment T<sub>3</sub> (row normal planting) (control) recorded the lowest fodder yield (8.74 q ha<sup>-1</sup>) of green gram being *at par* with T<sub>4</sub>, indicating yield reduction due to normal planting pattern. These results confirmed the findings of Singh and Kaushik (1987), Padhi *et al.* (2001) with respect to grain and fodder yields.

The data showed that the differences in harvest index of green gram due to planting pattern and fertility levels were non-significant.

The data on economics of different planting pattern and fertility levels of sweet corn and green gram clearly indicated that T<sub>8</sub> (Sweet

corn + green gram (2:2) in paired row of 45/90 cm) (100 % RFD of N in sweet corn) secured maximum net realization of Rs. 64395 ha<sup>-1</sup> with BCR of 5.63, followed by Rs. 60947 ha<sup>-1</sup> with BCR of 5.28 in T<sub>9</sub> (Sweet corn + green gram (2:2) in paired row of 45/90 cm) (125 % RFD of N in sweet corn). These results are in conformity with those reported by Padhi (2001) and Ghosh and Singh (1996).

**REFERENCES**

Akhtar, M. and Silva, J.A. 1999. Agronomic traits and productivity of sweet corn affected by nitrogen and inter-cropping. *Pakistan J. Soil Sci.*, **16** : 49-52.

Anonymous 2004. A little impetus needed. *The Hindu Survey of Indian Agric.*, pp 61-63.

Arya, M.P.S. and Saini, R.P. 1989. Effect of planting geometry on maize and soybean intercropping systems under rainfed conditions. *Indian J. Agron.* **34** : 322-24.

Dhingra, K. K., Dhillon, D. S., Grewal, D. S. and Sharma, K. 1991. Performance of maize and mungbean intercropping in different planting pattern and row orientations *Indian J. Agron.* **36** : 207-12.

Ghosh, P. K. and Singh, N. P. 1996. Production potential of summer legumes-maize sequence under varying levels of nitrogen. *Indian J. Agron.* **41** : 525-28.

Padhi, A.K. 2001. Effect of vegetable intercropping on productivity, economics and energetics of maize *Indian J Agron.* **46** : 204-10.

Padhi, A.K. and Panigrahi, R.K. 2006. Effect of intercropping and crop geometry on productivity, economics, energetic and soil fertility status of maize based intercropping systems. *Indian J. Agron.* **51** : 174-77.

Shivran, R.K. and Rana, K.S. 2003. Growth and yield of maize as influenced by cropping systems and moisture concentration practices under rainfed conditions. *Annals Agric. Res.*, **24** : 350-53.

Sharma, R. S. 1995. Performance of maize-legume intercropping systems under varying nitrogen. *JNKVV Res. J.* **27**: 5-9.

Singh, R. P. and Kaushik, M. K. 1987. Nitrogen economy in maize-legume intercropping system. *Annals Agric. Res.*, **8** 1 : 105-09.

Wiley, R.W. 1979. Intercropping its importance and research needs part-I competition and yield advantages. *Field Crop Abst.* **32**: 1-10