

## CONTENT

| Title   | Author   | Page No. |
|---|--|----------|
| <b>Production</b>   |  |          |
| Cloning and characterization of anthocyanin biosynthetic regulatory gene of purple sweet potato ( <i>Ipomoea batatas</i> L.)  | N. Choudhury, J. Tarafdar and P. P. Sur                                  | 01-16    |
| Nutrient scheduling for upland rice intercropped in coconut   | B. M. Suman, K. Prathapan, S. K Raj, N.V. Radhakrishnan and E. K. Syriac | 17-23    |
| Productivity, phenology and heat unit accumulation of wheat ( <i>Triticum aestivum</i> L.) as influenced by nitrogen management using green seeker                            | J. Kaur, H. Ram and S. Pal Singh   | 24-28    |
| Induction of flowering in mango cv. Himsagar  | S. Kundu, J. Mishra, P. Nandi, R. Swamy Sekhar and T. Adhikary           | 29-34    |
| Rejoinder of diverse organic growing media on morphological attributes of nine <i>Heliconia</i> species and varieties under West Bengal condition                             | M. Malakar, P. Acharyya and S. Biswas                                    | 35-44    |
| Hybrid maize (QPM) for nutritional security   | B. Subbulakshmi  | 45-49    |
| Effect of tree architecture and variety on growth and yield attributes of apple under UHDP  | K. K. Srivastava, D. Kumar, S. R. Singh, O. Chand and S. Pandey          | 50-55    |
| A review on developments in oil extraction from oilseeds  | K. Dhali and N. Basak  | 56-63    |
| Study on genetic variability and correlation for floral traits and yield in <i>Heliconia</i> genotypes  | M. Malakar, P. Acharyya and S. Biswas                                    | 64-72    |
| Performance of marker assisted back cross bread wheat ( <i>Triticum aestivum</i> L.) varieties in relation to sowing environment  | C. Kaur, H. Ram and G. S. Mavi   | 73-77    |
| Evaluation of some watermelon hybrids and cultivars under Lateritic Belt of West Bengal   | S. Mohanta and J. Mandal   | 78-82    |
| Extricate the impact of tactics of nitrogen source on development of <i>Lilium</i> cultivars  | M. A. Wani, I. T. Nazki, A. A. Sheikh, A. Din, T. A. Bhat and R. A. Wani | 83-91    |
| Effect of micronutrients on growth, yield and quality of strawberry ( <i>Fragaria</i> × <i>ananassa</i> Duch.) cv. Winter Dawn in the Gangetic Alluvial Region of West Bengal | T. Saha, B. Ghosh, S. Debnath and A. Bhattacharjee                       | 92-95    |

| <b>Title</b>   | <b>Author</b>   | <b>Page No.</b> |
|--|---|-----------------|
| Nitrogen indices of wheat sown with happy seeder with skipped row techniques as affected by split nitrogen application                               | R. K. Hundal and J. S. Kang                                       | 96-101          |
| Estimation of missing observations of agricultural experiments under split plot set up   | A. Dutta, A. Majumder, D. Nishad and S.K. Mandi                   | 102-107         |
| Response of aerobic rice to irrigation and nitrogen management in red and lateritic soil of West Bengal  | S. Duary and K. Pramanik  | 108-113         |
| Effect of crop combination and nutrient management on yield, nutrient uptake and economics of sweet corn based cropping system                       | B. Swain, L. M. Garnayak and S. Mangaraj                          | 114-120         |
| <b>Protection</b>  |   |                 |
| Influence of weed control techniques and establishment method on yield and economics of rainfed lowland rice   | J. Lhungdim, Y. S. Devi, K. N. Devi and Y. B. Chanu               | 121-126         |
| Soil microbes, nutrient removal, productivity and profitability of lentil ( <i>Lens culinaris</i> Medikus) influenced by weed management practices   | H. Kalita and R. Chakrabarty                                      | 127-133         |
| Dissipation of insecticides in okra fruits   | S. A. Pandey, P. Mishra and V. K. Koshta                          | 134-138         |
| Arsenic contamination through food-chain in Gangetic delta of West Bengal – looking beyond drinking water pollution menace                           | B. Sinha, K. Bhattacharyya and P. K. Giri                         | 139-144         |
| Ergonomical studies on single row power weeder for rice crop   | S. Pandey, A. K. Verma and A. Sirmour                             | 145-150         |
| Diversity and dynamics of weeds as influenced by flucetosulfuron 10 % WG in direct seeded rice and its residual effect on succeeding crop green gram | S. Sridhara, P. Gopakkali, T.B. Naik., S. Pradeep and H.D. Shilpa | 151-157         |
| Efficacy of herbicides against weed flora in onion ( <i>Allium cepa</i> L.)  | B. B. Sahoo and P. Tripathy                                       | 158-163         |
| Effect of herbicides on weed dynamics, phytotoxicity and seed yield of linseed ( <i>Linum usitatissimum</i> L.)                                      | C. S. Puhup and S. K. Dwivedi                                     | 164-169         |
| Evaluation of leguminous cover crop and banana bio-mat mulching for weed suppression and conservation of soil moisture and nutrient in guava orchard | A. Bhattacharjee and S. Debnath                                   | 170-177         |
| <b>Short Communication</b>   |   |                 |
| Constraints faced by the organic farmers of Manipur State, India   | A. Hanglem, S. Modak, D. Roy and P. K. Pal                        | 178-181         |
| Effect of bio-fertilizer and mulching on growth, yield and quality of cauliflower ( <i>Brassica oleracea</i> var. <i>botrytis</i> L.) in Punjab      | G. Singh and 'S. K. Singh   | 182-185         |

| <b>Title</b>   | <b>Author</b>   | <b>Page No.</b> |
|--|---|-----------------|
| Study on the performance of mungbean varieties in the New Alluvial Zone of West Bengal   | R. Mondal and K. Sengupta   | 186-191         |
| Yield, quality and soil fertility as influenced by <i>rabi</i> castor ( <i>Ricinus communis</i> L.) based intercropping system                     | B. K. Yadav, G. N. Patel, and S. K. Choudhary   | 192-194         |
| Performance of zero till seed drill in raising blackgram under rice fallow   | G. Amuthaselvi, V. Dhanushkodi and S. Eswaran   | 195-197         |
| Effect of moisture conservation on growth and yield of linseed under varying fertility levels  | A. K. Devedee, R. K. Singh, R. N. Meena and K. Choudhary                                  | 198-200         |
| Variation in seed production potential of China aster genotypes in the New Alluvial Zone of West Bengal  | A. Chakraborty, S. K. Bordolui, M. K. Mahato, R. Sadhukhan and D. J. M. S. N. K. Sri Veda | 201-204         |
| Effect of humic acid on the growth and yield of <i>rabi</i> pigeon pea [ <i>Cajanus cajan</i> (L.) Millsp] in the New Alluvial Zone of West Bengal | A. Nalia and K. Sengupta  | 205-208         |
| Influence of different doses of pyrazosulfuron-ethyl and establishment methods on the yield of lowland rice ( <i>Oryza sativa</i> L.)              | T. Malemnganbi and J. Lhungdim  | 209-212         |
| Rapid and consistent QuEChERS based RP-HPLC methodology for the detection of various insecticides in cabbage and cauliflower                       | R. Akoijam and R.S. Telem   | 213-217         |
| Diversity of weed flora in pineapple plantations of Kerala   | T. Girija and M. V. Menon   | 218-221         |