

## CONTENT

| Title   | Author  | Page No. |
|---|---|----------|
| A study on genetic variability for yield and its attributes in sacred basil ( <i>Ocimum tenuiflorum</i> L.)                                       | M. Pavan Gowda, A. V. D. Dorajeerao, M. Madhavi and D.R. Salomi Suneetha  | 01-06    |
| Evaluation of F <sub>4</sub> segregating population of bitter gourd ( <i>Momordica charantia</i> L.) for qualitative and quantitative traits      | P. Pradhan, P. Thipathy, G. S. Sahu, S. K. Dash, B. Pradhan, B. Tripathy, P. Sahu, N. J. Nayak, S. Saurav and S. Mishra   | 07-11    |
| Sustainable approach towards conversion of unproductive coconut palms into productive ones through neera production                               | Tokoyarin, A. Bandyopadhyay and D. K. Ghosh (LKN)   | 12-18    |
| Identification of sorghum parental lines with high phenotypic stability using GGE biplot analysis in Central India                                | U. Saxena, S. Reddy, M. K. Saxena and P. Kumari   | 19-24    |
| Growing mango: a profitable livelihood option for Bhagalpur farmers   | K. V. Vijay, S. M. Rahaman, M. K. Wadhvani, S. Kumari, M. Kumari, S. Kumar, F. Homa, S. Sengupta and S. L. Bairwa         | 25-27    |
| Impact of different mulch materials on growth and flowering of rose cv. <i>Mainu Parle</i>  | L. Tripathy, S. K. Dash and T. K. Giri  | 28-31    |
| Effect of lime, micronutrient and spraying schedule on growth and yield of broccoli variety <i>Palam Samridhi</i>                                 | S. Sourabh, P. Tripathy, G. S. Sahu, S. K. Dash, R. K. Nayak, P. Sahu, N. J. Nayak, B. Tripathy, P. Pradhan and S. Mishra | 32-38    |
| Perception of tribal farmers on fish culture practices in small water bodies: an exploratory study in Khowai district of Tripura, India           | S. P. Debbarma, A. Ghosh, B. Lahiri and P. K. Pandey  | 39-43    |
| Assessment of potato ( <i>Solanum tuberosum</i> L.) – onion ( <i>Allium cepa</i> L.) cropping sequence under lower Gangetic plains of West Bengal | S. K. Das and A. Chakraborty  | 44-47    |
| Prognostication of mulberry silk cocoon prices in Kaliachak (West Bengal) market  | Manjunatha G. R., R. Hunmily E, K. K. R. Patil, S. Afroz, Parmeshwarnaik J., D. Pandit and V. Sivaprasad                  | 48-53    |
| Effect of foliar spray on phenology and yield of Lentil sown on different dates   | V. Visha Kumar, P. Banerjee, R. Nath, K. Sengupta, Sarath Chandran M. A. and R. Kumar                                     | 54-58    |
| Climate smart agriculture for sustaining food production  | D. Nongmaithem, M. Apon, A. P. Singh and L. Tzudir  | 59-64    |
| Breeding for improved leaf yield and studies on combining ability in mulberry   | K. Suresh, Jalaja, S. Kumar, D. Chakravarty and V. Sivaprasad   | 65-71    |
| Seed priming and foliar nutrition studies on relay grass pea after winter rice in lower Gangetic plain  | P. Banerjee, V. Visha Kumari, R. Nath and P. Bandyopadhyay  | 72-78    |

| <b>Title</b>  | <b>Author</b>  | <b>Page No.</b> |
|---|--|-----------------|
| Phenotypic stability in brinjal genotypes   | S. Akhtar, Aakanksha, R. Kumari, S. S. Solankey and D. K. Baranwal | 79-86           |
| Standardization of integrated nutrient management for aromatic <i>Gobindabhog</i> rice in <i>Gangetic</i> alluvial region of West Bengal                  | D. Mahata, M. Ghosh, B. C. Patra, S. K. Pal and S. Banerjee        | 87-92           |
| Comparative seasonal influence of <i>Kuliana lime</i> for fruit and yield parameters under Bhubaneswar condition  | S. Mishra and D. K. Dash   | 93-99           |
| Effect of antitranspirants on rice ( <i>Oryza sativa</i> ) grown under submerged and water stressed condition in an inceptisol of Varanasi, Uttar Pradesh | R. Patel, P. Raha, Sri Laxmi and A. Paul                           | 100-106         |
| Quality assessment in association with yield attributes contributing improved yield in onion ( <i>Allium cepa</i> L.)                                     | S. Bal, T. K. Maity, A. B. Sharangi and A. Majumder                | 107-115         |
| Performance of medium duration rice varieties to varying levels of fertility and split application of nitrogen  | R. Pattnaik, M. R. Satapathy, K. Sethi and S. N. Jena              | 116-120         |
| Climate change perception and productivity of rice : the ground truth verification of two selected villages of Bankura district of West Bengal            | S. Mandal and T. K. Mondal   | 121-125         |
| Critical review on microbial perspective of jute retting  | S. Sinha Ray, S. Mukherjee, Md. N. Ali and B. Majumdar             | 126-130         |
| Enhancement in rice production as influenced by cropping systems and integrated nutrient management in New Alluvial Zone of West Bengal                   | A. Begam, P. Haldar, M. Ray, R. Debasharma and D. C. Roy           | 131-138         |
| Integrated management of rock bulrush ( <i>Schoenoplectus juncooides</i> ) in wet seeded rice   | F. Umkhulzum and Ameena. M   | 139-144         |
| Comparative physiology of salinity, drought and heavy metal stress during seed germination in ricebean [ <i>Vigna umbellata</i> (Thunb.) Ohwi and Ohashi] | K. Atta, S. Karmakar, D. Dutta, A. K. Pal and K. Jana              | 145-149         |
| Effect of drought mitigation strategies on growth and productivity of pigeonpea ( <i>Cajanus cajan</i> (L.) Millsp.) under rainfed conditions             | P. Venkata Rao, A. Subbarami Reddy and M. V. Ramana                | 150-153         |
| Physiological and biochemical parameters as a basis of screening salinity tolerance at seedling stage in Rice ( <i>Oryza sativa</i> L.)                   | Awaneet Kumar, S. K. Sarangi, A. B. Mandal                         | 154-159         |

| <b>Title</b>   | <b>Author</b>   | <b>Page No.</b> |
|--|---|-----------------|
| Evaluation of the efficacy of pre and post emergence herbicides to manage grassy and broad leaf weeds on mungbean( <i>Vignaradiata</i> L.) in Western Tigray | Zenawaigebregergis, Fischehabaraki and Gebreki Rosmaru      | 160-166         |
| Efficacy of pendimethalin against resistant little seed canary grass as affected by soil moisture and formulation  | M. Kaur, S. S. Punia, J. Singh and S. Gingh                 | 167-173         |
| Relevance of weed science in modern agriculture  | R. R. Upasani and S. Barla                                  | 174-177         |
| Identification and characterization of bacteria causing flacherie in mulberry silkworm, <i>Bombyx mori</i> L.  | K. Rahul, Moamongba, K. Saikia, M. Rbha and V. Sivaprasad   | 178-181         |
| Detection and molecular characterization of Begomovirus infecting some common weeds in West Bengal   | S. Chakraborty, T. Abing, S. Das and J. Tarafdar            | 182-191         |
| Efficacy of spirotetramat 11.01 + imidacloprid 11.01 SC against jassids, red mites and general predators in tomato   | L. C. Patel and A. Sarkar                                   | 192-200         |
| Bio-efficacy evaluation of oxyfluorfen 23.5% EC for controlling weeds in potato crop   | U. Bhattacharya, S. Sarkar and S. Dewanjee                  | 201-208         |
| Evaluation of synthetic antimicrobial peptides for the control of mulberry brown leaf spot disease incited by <i>Myrothecium roridum</i>                     | P. Makwana, S. Chattopadhyay and V. Sivaprasad              | 209-212         |
| Evaluation of glyphosate 41% SL to control weeds in tea at <i>Terai</i> region of West Bengal  | N. Meena Devi, C. K. Kundu, P. S. Bera, S. Panda and S. Das | 213-219         |
| Long term effect of fertilizer on weed dynamics  | Sheela Barla and R. R. Upasani                              | 220-224         |