

## CONTENT

| Title   | Author   | Page No. |
|---|--|----------|
| Factors affecting the production of major pulse crops in Madhya Pradesh<br>DOI : <a href="https://doi.org/10.22271/09746315.2022.v18.i3.1608">https://doi.org/10.22271/09746315.2022.v18.i3.1608</a>  | A. Singh and S. Singh  | 1-7      |
| Nitrogen mineralisation of green manure crops during <i>rabi</i> - a relative analysis<br>DOI : <a href="https://doi.org/10.22271/09746315.2022.v18.i3.1609">https://doi.org/10.22271/09746315.2022.v18.i3.1609</a>   | M. Sanjana, G. S. Madhu Bindu, B. Padmaja, M. Uma Devi and S. Triveni  | 8-12     |
| Efficient nutrient management for profitable quality <i>tossa</i> jute ( <i>Corchorus olitorius</i> L.) seed production in the intensively jute fibre growing southern Bengal region<br>DOI : <a href="https://doi.org/10.22271/09746315.2022.v18.i3.1610">https://doi.org/10.22271/09746315.2022.v18.i3.1610</a> | S. Sarkar, M. Behera, A. Bera and S. K. Sarkar   | 13-18    |
| Assessment of forage berseem ( <i>Trifolium alexandrinum</i> L.) for productivity and profitability under varying seed rates and phosphorus fertilization<br>DOI : <a href="https://doi.org/10.22271/09746315.2022.v18.i3.1611">https://doi.org/10.22271/09746315.2022.v18.i3.1611</a>                            | M. Arif, A. Kumar, R. Pourouchottamane, D. L. Gupta and B. Rai   | 19-25    |
| Integrated nutrient management for improving growth and yield of cowpea<br>DOI : <a href="https://doi.org/10.22271/09746315.2022.v18.i3.1612">https://doi.org/10.22271/09746315.2022.v18.i3.1612</a>  | K. Pushpa, N. Pruthviraj and M. G. Vinay   | 26-30    |
| Variation in yield and yield attributes of hybrid mustard ( <i>Brassica juncea</i> M.) influenced by different spacing and fertilizer levels<br>DOI : <a href="https://doi.org/10.22271/09746315.2022.v18.i3.1613">https://doi.org/10.22271/09746315.2022.v18.i3.1613</a>   | A. Mala, S. K. Singh, R. Nath, A. John, P. Kumar and A. Bera   | 31-37    |
| Yield intensification of spring mungbean ( <i>Vigna radiata</i> L.) through exogenous foliar nutrition<br>DOI : <a href="https://doi.org/10.22271/09746315.2022.v18.i3.1614">https://doi.org/10.22271/09746315.2022.v18.i3.1614</a>   | T. Mahato, K. Murmu, A. Sarkar, S. Maji and P. Das   | 38-43    |
| Quality and suitability of fertilised <i>rabi</i> legume crops for green manuring<br>DOI : <a href="https://doi.org/10.22271/09746315.2022.v18.i3.1615">https://doi.org/10.22271/09746315.2022.v18.i3.1615</a>  | A. Chendra Babu Naidu, G. S. Madhu Bindu, M. Malla Reddy and M. Uma Devi   | 44-50    |
| Evaluation of AB-DTPA for multinutrient extraction from cardamom and pepper plantations in Idukki district of Kerala<br>DOI : <a href="https://doi.org/10.22271/09746315.2022.v18.i3.1616">https://doi.org/10.22271/09746315.2022.v18.i3.1616</a>   | V. I. Beena, P. Geetha, P. S. Philip, P. Sureshkumar, Amrutha S. Ajayan, Sam K. Issac, R. S. Sooraj and A. K. Shukla | 51-56    |
| Optimization of magnesium and boron dosage in cardamom grown in Idukki district of Kerala<br>DOI : <a href="https://doi.org/10.22271/09746315.2022.v18.i3.1617">https://doi.org/10.22271/09746315.2022.v18.i3.1617</a>  | V. I. Beena, P. Geetha, P. S. Philip, P. Sureshkumar, Amrutha S. Ajayan, R. S. Sooraj, Sam K. Issac and A. K. Shukla | 57-63    |
| Juxtaposition of morpho-qualitative traits of mango varieties grown in Mohanpur, West Bengal and Paria, Gujarat<br>DOI : <a href="https://doi.org/10.22271/09746315.2022.v18.i3.1618">https://doi.org/10.22271/09746315.2022.v18.i3.1618</a>  | K. Chakraborti, N. I. Shah and P. Patil  | 64-70    |

| Title  | Author   | Page No. |
|--|--|----------|
| Quality and green fodder productivity of forage maize ( <i>Zea mays</i> L.) as influenced by different seed priming techniques under rainfed situation<br>DOI : <a href="#">KWWRURUJQ</a> <a href="#">YYUQ</a> | K. Jana, A. Sarkar, R. Mondal and R. K. Agrawal  | 71-77    |
| Evaluation of glyphosate ammonium 79.2% SG for weed dry weight, crop phytotoxicity, yield, available NPK and economics of tea ( <i>Camellia sinensis</i> L.)<br>DOI :  | S. Tanha, P. Chopra, S. S. Rana, G. Singh and N. Kaur                                    | 78-85    |
| Effect of sodium para nitrophenolate 0.3% SL on growth and yield of chilli ( <i>Capsicum annuum</i> L.) in Inceptisols<br>DOI :  | S. K. Das, S. Samui and P. Nandy   | 86-91    |
| Weed dynamics as influenced by drip irrigation and mulching in guava<br>DOI :  | R. Jat, V.P. Singh, S. K. Guru, J. S. Shivran and K. Adhikary                            | 92-99    |
| Influence of organic weed management practices on phyto-sociology and diversity of weeds in maize-pea cropping system<br>DOI :   | G. Hetta, S.S. Rana, S. Kumar and M. Chakraborty   | 100-111  |
| Efficacy of module based integrated approach to combat major insect pests infesting summer green gram<br>DOI :   | A. Banerjee and S. Ray   | 112-119  |
| Bio-efficacy evaluation of insecticides against aphids infesting potato<br>DOI :   | S. Naik, A. K. Nayak, A. Sasmal and A. Mishra  | 120-127  |
| Integrated management of <i>Alternaria</i> blight caused by <i>Alternaria</i> spp. in rapeseed ( <i>Brassica rapa</i> var. toria) under field conditions<br>DOI :  | R. Chakrabarty, H. Kalita, M. Chakravarty, M. Basumutary, B. Bhattacharyya and B. Medhi  | 128-135  |
| Morpho-biochemical and molecular insights of fibre quality in jute: A review update<br>DOI :   | S. Bandyopadhyay, S. Datta and Md. N. Ali  | 136-151  |
| Stability investigation for seed yield and component characteristics in grain amaranth [ <i>Amaranthus hypochondriacus</i> L.] under different environments<br>DOI :   | P. M. Savaliya, N. N. Prajapati, R. S. Solanki, N. V. Soni, M. S. Patel and K. K. Tiwari | 152-157  |
| EMS induced mutagenesis in green gram for small seeded bruchid resistant genotype<br>DOI :   | A. Sen, A. P. Singh, S. Sarkar and S. Bhattacharyya                                      | 158-164  |

| <b>Title</b>   | <b>Author</b>   | <b>Page No.</b> |
|--|---|-----------------|
| Genetic diversity studies in muskmelon using microsatellite markers<br>DOI :   | G. Indrāja, S. Syed,<br>B. T. Priya and Rajesh  | 165-171         |
| Genetic variability studies of garlic ( <i>Allium sativum</i> L.) germplasm in Terai region of West Bengal<br>DOI :  | Y. R. Reddy, P. S. Medda,<br>B. Roy and S. Datta  | 172-180         |
| Comparative study on physico-chemical properties of F <sub>1</sub> hybrids and its parent of guava ( <i>Psidium guajava</i> L.) in Terai Agro-Climatic Zone of West Bengal<br>DOI :                                  | Z. W. Sherpa, A. Chakraborty, N. Bhowmick,<br>A. N. Dey, A. Sarkar and S. Khalko  | 181-184         |
| Evaluation of physico-chemical attributes and postharvest qualities of Himsagar mango with the exogenous application of growth regulators, anti-oxidant and geotextile mulch<br>DOI :                                | S. Dey, S. Datta, M. Alam and , P. Datta  | 185-189         |
| The efficacy of varying length of cladode cutting and IBA concentration on root and shoot growth in dragon fruit ( <i>Hylocereus costaricensis</i> [F.A.C. Weber] Britton and Rose.) cv. Royal Moroccan Red<br>DOI : | S. Chhetri, M. Alam, M. A. Hasan and<br>A. Lama   | 190-194         |
| Effect of accelerated ageing and dry dressing with botanicals on seed quality of soybean<br>DOI :  | V. R. Hridya, S. Lakshmi and S. Ambika  | 195-199         |
| Modelling and forecasting of electricity consumption used in agriculture purpose in India<br>DOI :   | R. K. Tiwari P. Mishra, B. Kumari,<br>A. M. G. Al Khatib, S. Yadav, M. Abotaleb,<br>S. Ray and M. Kumari                                    | 200-205         |
| Popularization of improved short duration rice variety Telangana Sona (RNR 15048) through frontline demonstrations in Nalgonda District, Telangana<br>DOI :  | M. Shankar, M.A. Aariff Khan, T. Bharath,<br>S. Pallavi, T. Himabindu, R.V.T. Balazzi Naaiik,<br>K. Sumalini, V. R. Naik And M. Shankaraiah | 206-212         |
| A study on entrepreneurial behavior of ericulture farmers in the Charaideo district of Assam<br>DOI :  | M. Marak, Th. A. Singha and E. Deori  | 213-217         |
| Attributes of CRIJAF Nail Weeder as perceived by the jute growers in West Bengal<br>DOI :  | D. Chatterjee, S. Mondal and D. Basu  | 218-222         |

| <b>Title</b>  | <b>Author</b>   | <b>Page No.</b> |
|---|---|-----------------|
| An economic impact of conservation agriculture on small and marginal farms in West Bengal, India: An assessment with DID method<br>DOI :  | S. Chatterjee, M. Hasan and R. Chakraborty  | 223-237         |
| An economic impact of lac based livelihood on tribal community of West Bengal, India<br>DOI :   | B. C. Mahato, S. De and S. Chatterjee   | 238-244         |
| Development of zonal wheat yield forecast models through Principal Component Analysis<br>DOI :  | Chetna, M. Devi, J. Mohammed,<br>Y. S. Raghav and P. Mishra                                 | 245-250         |
| Effect of crop combinations on nut equivalent yield and economics of coconut based cropping system<br>DOI :   | Namitha V.V., S. K. Raj, S. Pillai P.,<br>K. Prathapan, N. V. Radhakrishnan and<br>D. Jacob | 251-255         |
| Response to foliar application of different nutrient on lentil ( <i>Lens culinaris</i> Medik.) as relay crop under rice assessed in terms of performance of yield and economics in the Eastern Region of India<br>DOI : | A. Hoque, H. Singh, R. K. Srivastava<br>and A. Barman                                       | 256-259         |
| Influence of different nutrient management schedules on physiological attributes of organic cabbage ( <i>Brassica oleracea</i> var. <i>capitata</i> L.)<br>DOI :  | Sindhu V. and R. Chatterjee   | 260-263         |
| Evaluation of superior Darjeeling mandarin genotypes/ clones from Sikkim and Darjeeling hills<br>DOI :  | N. Gurung., S. Singh, D. Barman,<br>S. Sarkar, B. K. Singh and N. Prasad                    | 264-267         |
| Influence of different companion crops on major sucking pests in the brinjal ecosystem<br>DOI :   | M. I. Zaman, B. Mondal,<br>A. K. Mukhopadhyay and L. C. Patel                               | 268-272         |
| Estimation of cost of cultivation, profitability in different farm sizes as well as growth and instability of mustard in West Bengal<br>DOI :   | A. Roy, S. Sarkar and B. Bera   | 273-277         |