



## The effect of sarcotesta and time of sowing on seed germination of papaya (*Carica papaya* L.) cv. GJP-1

K.S. JOTAVA, D. R. KANZARIA AND \*S. MISHRA

Department of Horticulture, College of Agriculture, Junagadh Agricultural University, Junagadh, Gujarat, India

<sup>1</sup>Department of Horticulture, Institute of Agricultural Sciences, Sikhya 'O' Anusandhan University, Bhubaneswar, Odisha, India

Received : 24.12.2022 ; Revised : 13.01.2023 ; Accepted : 17.02.2023

DOI: <https://doi.org/10.22271/09746315.2023.v19.i1.1664>

### ABSTRACT

Slow and asynchronous germination of papaya seeds partly might be associated with the occurrence of various inhibitors in the gelatinous sarcotesta and seed coat. Similarly, the sowing of papaya seeds at different times affects the germination of papaya seeds. Hence, this experiment was undertaken at JAU, Gujarat to evaluate the influence of both the parameters viz., sarcotesta and time of sowing on seed germination of papaya cv. GJP-1. The removal of sarcotesta by hand rubbing was done and after evaluating the better treatment among the effect of sarcotesta, the seeds were shown in the polybag containing the media mixture of soil: sand: FYM in 1:1:1 ratio. Among the treatment effect of sarcotesta the seeds without sarcotesta executed the highest germination percentage (81.03%), survival percentage (98.29%) and least days for germination (14.28). Among the different dates of sowing, the early germination of papaya seeds (8.28 days) was found on 1<sup>st</sup> July ( $T_{1j}$ ) whereas, the highest germination and survival percentage (96.66 and 99.28 %) were observed in seeds sown on 16<sup>th</sup> June ( $T_{12}$ ) and 1<sup>st</sup> June ( $T_{11}$ ), respectively. Hence, it could be concluded that for obtaining good quality papaya seedlings, the seeds (cv. GJP-1) should be soaked for 24 hours in the water, followed by the removal of sarcotesta by hand rubbing and seeds should be sown during June to July months of the year for getting higher germination and survival percentage.

**Keywords:** Germination, GJP-1, papaya, sarcotesta, survival, time of sowing

Papaya (*Carica papaya* L.) also known as papaw or pawpaw belongs to the family Caricaceae with chromosome number 18. Accelerated demand for fresh fruit leads to an increase in papaya acreage. As a result of its quick and heavy yielding potential, this crop is grown around the tropics and subtropics of the World (Reddy and Gowda, 2014) producing a favourable Cost-Benefit ratio (Sharma and Zote, 2010). It is abundant in carbohydrates, vitamins and minerals (Ram, 2005). In recent days, farmers are getting interested in growing papaya as the main crop because it gives fruiting throughout the year and is regularly in demand. Papaya is commercially propagated through its seeds, which are protected by a gelatinous sarcotesta. Papaya seeds have been found to contain inhibitors, primarily phenolic compounds, in the sarcotesta and the seed coat, which contribute to slow and asynchronous germination (Chow and Lin, 1991; Reyes *et al.*, 1980) and affect the subsequent seedling emergence (Malo and Campbell, 2001). The powerful germination inhibitors in the fruit flesh prevent the inhibitor from travelling from the ovary flesh to the seed coat. Also, the papaya seed's sarcotesta/aril may hinder germination by blocking oxygen and water from reaching the embryo (Yahiro, 1979; Angeline and Ouma, 2008; Rodriguez *et al.*, 2019; Sharma *et al.*,

2021). So, it is recommended that, after harvesting, the seeds be gently rubbed with sand and thoroughly washed under running water to get rid of the gelatinous sarcotesta before drying and storage. Soaking papaya seeds in water for 24 hours prior to planting helps in better germination (Riley, 1981).

Seed germination and plant development are profoundly influenced by environmental conditions. It has been demonstrated that irregular shifts in climatic parameters have an impact on the success of seed germination and the development of seedlings. Variations in germination and seedling growth parameters result from the sowing of seeds at different times. Climatic parameters like temperature, humidity, bright sunshine hours, rainfall *etc.*, are important factors for nursery growers for successful seedling preparation. One of the most influential aspects of seed germination is temperature. Very high or low temperatures may inversely affect the process of germination. Papayas thrive in temperatures between 26 and 30 degrees Celsius during their early stages of growth in the nursery (Ram 1996; Ram and Ray, 1992). The relative humidity of the atmosphere has a direct influence on nursery raising. The optimum relative humidity is necessary to boost seed germination and seedling growth. The bright

Email: [subhrajyoti.ss235@gmail.com](mailto:subhrajyoti.ss235@gmail.com)

How to cite : Jotava, K.S., Kanzaria, D. R. and Mishra, S. 2023. The effect of sarcotesta and time of sowing on seed germination of papaya (*Carica papaya* L.) cv. GJP-1. *J. Crop and Weed*, 19 (1): 84-87.

sunshine hour is the key part of the growth of any plant as plants require bright sunshine to make their food for running the normal lifecycle. Inadequate sunshine hours cause the yellowing of plants and will result in uneven and suppressed growth of seedlings. Rainfall will improve the microclimate that is suitable for seed germination and seedling growth. But the continuous and heavy rainfall will decrease the BSS, increase humidity in the atmosphere and will make the soil compact which may hinder seedling growth. Thus, different climatic parameters have varied influences on the seed germination of papaya. So, it is important to study the right time of sowing for papaya for getting higher germination success and good quality seedlings as now-a-days farmers are interested to grow gynodioecious varieties which are very costly. For farmers, every seed is worthy and important. The farmers do not have infrastructure facilities, like playhouses, greenhouses *etc.*, for raising seedlings under protected conditions. They grow the seedlings under the shed or under field conditions. The farmers with small land holdings have poor purchasing power and sowing seeds with the wrong method and time will cause economical loss.

#### MATERIALS AND METHODS

The present investigation was carried out at Fruit Research Station, Lal Baugh, Junagadh Agricultural University, Junagadh, Gujarat. The treatments comprised of  $S_1$  (with sarcotesta) and  $S_2$  (without sarcotesta) and  $T_1$ -1<sup>st</sup> January,  $T_2$ -16<sup>th</sup> January,  $T_3$ -1<sup>st</sup> February,  $T_4$ - 16<sup>th</sup> February,  $T_5$ -1<sup>st</sup> March,  $T_6$ -16<sup>th</sup> March,  $T_7$ -1<sup>st</sup> April,  $T_8$ -16<sup>th</sup> April,  $T_9$ -1<sup>st</sup> May,  $T_{10}$ -16<sup>th</sup> May,  $T_{11}$ -1<sup>st</sup> June,  $T_{12}$ -16<sup>th</sup> June and  $T_{13}$ -1<sup>st</sup> July as time of seed sowing. A Completely Randomized Design (CRD) with three replicates was used to plan the experiment. The papaya seed of variety GJP-1 was procured from Fruit Research Station, Madhadi Baug, Junagadh Agricultural University, Junagadh. Seeds were dipped in water for 24 hours and then gently rubbed with hands for the removal of sarcotesta. This was done precisely and with gentle hands so that it removes sarcotesta only and does not damage the seeds. The seeds were washed once or twice and immediately Carbendazim 2% powder was mixed to avoid any fungal infection. After mixing the Carbendazim seeds were spread on the muslin cloth to avoid sticking off with each other for one hour in the shed. Then seed sowing was done with three seeds sown in each polybag at 1.0 cm depth in a triangle manner. Seed sowing was done from January to July at 15 days intervals in black polybags containing soil: sand: FYM in 1:1:1 ratio.

#### RESULTS AND DISCUSSION

The perusal of data presented in Table 1 revealed that sarcotesta executed a significant influence on the days taken to germinate the papaya seeds. Significantly the early germination of papaya seeds (14.28 days) was found in seeds without sarcotesta ( $S_2$ ) and significantly maximum days (17.19 days) required for germination in the case of seeds with sarcotesta ( $S_1$ ). The better seed germination in papaya seeds without sarcotesta might be attributed to the fact that the removal of inhibitors helps in synchronous germination (Singh and Singh, 1981; Reyes *et al.*, 1980; Perez *et al.*, 1980; Tokuhisa *et al.*, 2007; Sarana *et al.*, 2016). Significantly the highest percentage (81.03%) of germination was observed in seeds without sarcotesta ( $S_2$ ) and significantly the lowest germination percentage (73.17%) was recorded in seeds with sarcotesta ( $S_1$ ). Significantly the highest survival percentage (98.29%) was observed in seeds without sarcotesta ( $S_2$ ) and significantly the lowest survival percentage (95.86%) was recorded in seeds with sarcotesta ( $S_1$ ). The sarcotesta on papaya seed may reduce germination by blocking oxygen infiltration into the seed, which may explain why seeds without sarcotesta have a higher germination rate (Yahiro, 1979; Angeline and Ouma, 2008). Aril acts as a physical barrier, restricting the flow of water and gases into the seeds; combined with the influence of phytohormones, this prevents germination, induces dormancy, stunts embryonic development, and reduces the overall germination rate (Rodriguez *et al.*, 2019). Some byproducts of anaerobic metabolism, as reported by Kranner *et al.* (2010), can hinder respiratory transport, setting off a chain reaction that ultimately resulted in the loss of cell function and kills the seed. The survival rate was lowest for seedlings grown from seeds infected with sarcotesta, possibly because this parasite interferes with germination and subsequent plant growth (Sangakara, 1995; Okeyo and Ouma, 2008).

Sowing papaya seeds at different times has a significant effect on the germination parameters of papaya seeds (Table 2). The early germination of papaya seeds (8.28 days) was found in treatment on 1<sup>st</sup> July ( $T_{13}$ ). The highest germination and survival percentage (96.66 and 99.28%) were observed in seeds sown on 16<sup>th</sup> June ( $T_{12}$ ) and 1<sup>st</sup> June ( $T_{11}$ ) respectively. The seeds shown on 1<sup>st</sup> July give early germination. It might be due to the fact that environmental conditions play a key role in the germination of seeds in a nursery. The favourable conditions of the month of July provide an adequate environment for the early germination of papaya seeds. The maximum germination percentage on 16<sup>th</sup> June might be attributed to the microclimate of that month like average temperature (29.58°C), humidity (70.5 %) and rainfall (101.66 mm with an average of

**Table 1: Effect of sarcotesta on days taken to germinate the papaya seeds, germination percentage and survival percentage of papaya seedlings**

Treatments	Days to germination	Germination percentage	Survival percentage
S <sub>1</sub> : With sarcotesta	17.19	73.17	95.86
S <sub>2</sub> : Without sarcotesta	14.28	81.03	98.29
<b>SEm (±)</b>	<b>0.108</b>	<b>0.558</b>	<b>0.398</b>
<b>LSD (0.05)</b>	<b>0.31</b>	<b>1.59</b>	<b>1.13</b>

**Table 2: Effect of time of sowing on days taken to germinate the papaya seeds, germination percentage and survival percentage of papaya seedlings**

Treatments	Days to germination	Germination percentage	Survival percentage
T <sub>1</sub> : 1 <sup>st</sup> January	23.83	59.79	94.39
T <sub>2</sub> : 16 <sup>th</sup> January	23.52	57.76	94.63
T <sub>3</sub> : 1 <sup>st</sup> February	20.20	66.39	95.37
T <sub>4</sub> : 16 <sup>th</sup> February	17.75	66.82	95.09
T <sub>5</sub> : 1 <sup>st</sup> March	18.10	69.28	98.86
T <sub>6</sub> : 16 <sup>th</sup> March	16.70	77.11	99.14
T <sub>7</sub> : 1 <sup>st</sup> April	16.18	67.94	99.06
T <sub>8</sub> : 16 <sup>th</sup> April	14.52	73.50	98.47
T <sub>9</sub> : 1 <sup>st</sup> May	12.95	82.98	98.99
T <sub>10</sub> : 16 <sup>th</sup> May	11.23	94.06	98.42
T <sub>11</sub> : 1 <sup>st</sup> June	11.55	93.45	99.28
T <sub>12</sub> : 16 <sup>th</sup> June	9.75	96.66	98.64
T <sub>13</sub> : 1 <sup>st</sup> July	8.28	96.55	91.65
<b>SEm(±)</b>	<b>0.275</b>	<b>1.424</b>	<b>1.015</b>
<b>LSD (0.05)</b>	<b>0.78</b>	<b>4.04</b>	<b>2.88</b>

3.5 days). The reduction in the germination of seeds sown during January and February might be due to low temperatures (Cheema and Dani, 1930; Naik, 1949). The maximum survival percentage on 1<sup>st</sup> June is might be due to the environmental conditions of that time (average temperature 30.66°C, average relative humidity 57 %, average rainfall 75.47 mm and average three rainy days). These environmental conditions are relatively congenial for the papaya seedling growth which got more survival percentage.

#### CONCLUSION

On the basis of the results obtained, it can be comprehended that for the preparation of papaya seedlings in the nursery, the seeds of cv. GJP-1, should be soaked for 24 hours in water and after removal of sarcotesta by hand rubbing the seeds should be sown from 16<sup>th</sup> May to 1<sup>st</sup> July for getting higher germination and survival percentage.

#### ACKNOWLEDGEMENT

The authors would like to thank the Department of Horticulture, Junagadh Agricultural University, Gujarat, for supporting this research.

#### REFERENCES

- Angeline, O. and Ouma, G. 2008. Effect of washing and media on the germination of papaya seeds. *J. Agric. Biol.Sci.*, **3**(1): 8-11.
- Cheema, G.S. and Dani, P.G. 1930. Bombay Agricultural Bull., No. 162. pp. 325 in Fruits of India: Tropical and Subtropical, 1<sup>st</sup> Edition (T. K. Bose, ed.) Naya Prokash, Calcutta 100006.
- Chow, Y.J. and Lin, C.H. 1991. P- Hydroxy benzoic acid as the major phenolic germination inhibitor of papaya seed. *Seed Sci. Technol.*, **19**: 167-174.
- Kranner, I., Minibayeva, F.V., Beckett, R.P. and Seal, C.E. 2010. What is stress? Concepts, definitions and applications in seed science. *New Phytologist*, **188**(3): 655- 673.
- Malo, E.A. and Campbell, C.W. 2001. Papaya: a potential annual crop under middle Georgia conditions. *J. Amer. Soc. Hortic. Sci.*, **115**(3): 426-438.
- Naik, K.G. 1949. South Indian Fruit and Their Culture. Varadachary and Co. Madras.
- Okeyo A. and Ouma, G. 2008. Effects of washing and media on the germination of papaya (*C. papaya*) seeds. *Agricultura Tropica et Subtropica*, **41**(1): 21-26.

- Perez, A., Reyes, M.N. and Cuevas, J. 1980. Germination of two papaya varieties: effect of seed aeration, K- treatment, removing of the sarcotesta, high temperature, soaking in distilled water and age of seeds. *J. Agric. Univ. Puerto Rico*, **64**: 173-180.
- Ram, M. 2005. Papaya. Indian Council of Agricultural Research, New Delhi. pp. 142-143.
- Ram, M. and Ray, P.K. 1992. Influence of fruiting season on seed production of papaya in north bihar. Abstracts VIII. All India seed seminar on recent advances and future strategies in seed science and technology, Nauni, Solan, June 9-11, 1992. H.P., India, pp. 27.
- Reddy, P.V.K. and Gowda, V.N. 2014. Influence of greenhouse cultivation on fruit quality of 'Red Lady' papaya. *Acta Hort.*, **1024**: 10.
- Reyes, M.N., Perez, A. and Cuevas, J. 1980. Detecting, endogenous growth regulators on the sarcotesta, endosperm and embryo by paper chromatography on fresh and old seeds of two papaya varieties. *J. Agric. Univ. Puerto Rico*, **64**: 164-172.
- Riley, J.M. 1981. Growing rare fruit from seed. California rare fruit grower's yearbook, **13**: 1-47.
- Rodriguez, S., Vargas, I., Hijuelo, A., Loumeto, F., Silva, J., Perez, J., Arias, Q., Fonseca, Y., Gomez, Y., Baldoquin, M. and Oliva, D. 2019. Analysis of the effect of scarification process on papaya (*Carica papaya* L.) seeds germination. *Intechopen*, 88012.
- Sangakara, U.R. 1995. Influence of seed ripeness, sarcotesta, drying and storage on germinability of papaya (*Carica papaya* L.) seed. *Pertanika J. Trop. Agric. Sci.*, **18**(3): 193-199.
- Sarana, P.L., Choudhary, R., Solanki, I.S. and Ercisli, S. 2016. Influence of temperature on seed germination in papaya under subtropical conditions of India. *Erwerbs- Obstbau*, DOI10.1007/s10341-016-0275-9.
- Sharma, P., Yadav, R., Jain, M. and Bhatishwar, C., 2021. Growing media and cow urine influence the seed germination and seedling growth of Papaya (*Carica papaya* L.). *J. Crop Weed*, **17**(3):253-259.
- Singh, R.M. and Singh, I.D. 1981. Effects of methods and duration of storage on seed germination and seedling vigour in papaya. *Seed Res.*, **9**: 67-72.
- Tokuhisha, D., Dias, D.C.F.S., Alvarenga, E.M., Hilst, P.C. and Demuner, A.J. 2007. Phenolic compound inhibitors in papaya seeds (*Carica papaya* L.). *Revista Brasileira de Sementes*, **29**(3): 180-188.
- Yahiro, M. 1979. Effect of seed pretreatments on the promotion of germination in papaya (*Carica papaya* L.). Memories of the faculty of agriculture, *Kago Shima Univ.*, **15**: 49-54.