



## Effect of plant growth promoting micro-organisms on the performance of strawberry

G. M. SANTHOSHKUMAR AND \*N. BHOWMICK

Department of Pomology and Post-Harvest Technology, Uttar Banga Krishi Viswavidyalaya, Pundibari, Cooch Behar, West Bengal-736165, India

Received : 05.12.2022 ; Revised : 24.01.2023 ; Accepted : 17.02.2023

DOI: <https://doi.org/10.22271/09746315.2023.v19.i1.1668>

### ABSTRACT

The present investigation was conducted at Uttar Banga Krishi Viswavidyalaya, Pundibari, West Bengal during 2020-21 and 2021-22 to assess the influence of plant growth promoting micro-organisms on the performance of strawberry. It was laid out in Randomized Block Design with eight treatments and three replications. The results revealed that, highest plant height (6.64 cm) was recorded with T<sub>8</sub> (Azotobacter + PSB + Trichoderma) at 30 days after planting, fresh weight of root (9.57 g), dry weight of roots (2.78 g) and number of crowns/plant (9.15). Whereas, T<sub>6</sub> (Azotobacter + Trichoderma) produced maximum number of leaves plant<sup>-1</sup> (5.50), shoot length (6.63 cm) at 30 days after planting, fresh weight of shoot (27.99 g) and dry weight of shoot (6.24 g) as compared to the control (T<sub>1</sub>). Estimated fruit yield (27.29 t ha<sup>-1</sup>) was recorded highest with T<sub>8</sub> (Azotobacter + PSB + Trichoderma), followed by T<sub>6</sub> (26.50 t ha<sup>-1</sup>) treated plants, and minimum in control T<sub>1</sub> (21.27 t ha<sup>-1</sup>).

**Keywords:** Azotobacter; fruit yield, growth, plant growth promoting micro-organisms, strawberry.

Strawberry (*Fragaria × ananassa* Dutch) is a fruit crop of major significance worldwide and has a basic chromosome number x=7. It belongs to the family Rosaceae (Ruiz *et al.*, 2011). Strawberry has drawn most growers' attention due to its nutrient value and importance in human diet (Giovannoni, 2004). The area under strawberry cultivation is increasing because of its diverse ecological state and cultivars having adaptability in diverse environmental conditions. In India, the crop is grown in West Bengal, Uttar Pradesh, Rajasthan, Punjab, Meghalaya, Mizoram, Maharashtra, Himachal Pradesh, Haryana, and Delhi. Sub-tropical regions in Jammu have also the potential to raise the crop under irrigated condition.

Strawberry grown best in a temperate climate and some of the varieties grown in milder subtropical climate. It is a short-day plant and fruits are highly perishable. Generally, the farmers are using chemical fertilizers to increase strawberry yield, resulted to affects the human body (Islam *et al.*, 2005). To overcome this problem, intensified use of beneficial micro-organisms and balanced integrated nutrient management improves the growth, flowering, and crop yield (Fernandes *et al.*, 2012; Cvijanovic *et al.*, 2007; Bhagat and Panigrahi, 2020). Beneficial micro-organisms enhances growth of plant and improves the soil health. Bio-fertilizers such as Arbuscular Mycorrhiza Fungi, *Trichoderma harzianum*, Phosphorous Solubilizing Bacteria, along with organic manures like vermicompost improves yield, mainly in reclaimed soils by overcoming salt, drought,

and some pathogenic stress in addition to reducing the applied fertilizers and enhancing the macro and micro elements availability (Estiken *et al.*, 2005; Balakrishna *et al.*, 2005; Mia *et al.*, 2005 and Sabarad *et al.*, 2004). Bio-fertilizers improves the physical properties of soil, enrich the soil nutrients, and enhance their effective plant absorption (Hazarika *et al.*, 2014).

Generally, Bio-fertilizers containing living micro-organisms, which improves the nutrient uptake capacity of the plant. Use of nitrogen fixing bacteria and phosphorous solubilizing bacteria supplying the essential nutrients like nitrogen and phosphorous by plant roots (Singh *et al.*, 2011). Biofertilizers keep the soil environment rich, increasing nutrient availability in soils and promote plant growth. Biofertilizers are preferred over synthetic fertilizers due to excess use of chemical which are harmful to the humans and biofertilizers do not reduce the soil fertility while continuous use (Umar *et al.*, 2009; Kumari *et al.*, 2018). Therefore, the present experiment has been made to study the influence of plant growth promoting micro-organisms on the performance of strawberry.

### MATERIALS AND METHODS

The field experiment was conducted at Uttar Banga Krishi Viswavidyalaya, Pundibari, Cooch Behar, West Bengal, India during 2020-21 and 2021-22, which is located at 42 meters above mean sea level, longitude 89°23'5'' E and latitude 26°19'86'' N, under the Sub-Himalayan Terai region of West Bengal. Chemical

Email: [nilesh@ubkv.ac.in](mailto:nilesh@ubkv.ac.in)

How to cite : Santhoshkumar, G. M. and Bhowmick, N. 2023. Effect of plant growth promoting micro-organisms on the performance of strawberry. *J. Crop and Weed*. 19 (1): 106-109.

fertilizers such as nitrogen (100 kg ha<sup>-1</sup>), phosphorous (80 kg ha<sup>-1</sup>), potassium (100 kg ha<sup>-1</sup>) and FYM (5 t ha<sup>-1</sup>) were applied into the soil one week before planting. Planting was performed at a distance of 25 × 50 cm and each bed was 2 × 1m apart. The strawberry planting materials were collected from a reputed nursery and the plant growth promoting micro-organisms were collected from Bio-Control Laboratory, Dept. of Plant Pathology, UBKV, Pundibari. Eight treatments and replicated thrice in a Randomized Block Design (RBD). Treatment details such as T<sub>1</sub> (Control), T<sub>2</sub> (*Azotobacter*), T<sub>3</sub> (Phosphate Solubilizing Bacteria), T<sub>4</sub> (*Trichoderma*), T<sub>5</sub> (*Azotobacter* + PSB), T<sub>6</sub> (*Azotobacter* + *Trichoderma*), T<sub>7</sub> (PSB + *Trichoderma*), T<sub>8</sub> (*Azotobacter* + PSB + *Trichoderma*) applied by soil drenching method at 15 days after planting (DAP). The parameters were subjected to analysed with the help of OPSTAT statistical package.

## RESULTS AND DISCUSSION

### Growth attributes

There was a significant enhancement in all growth attributes of strawberry due to use of beneficial micro-organisms (Table 1). The highest plant height (6.64 cm) was recorded in T<sub>8</sub> at 30 days after planting (DAP) followed by T<sub>6</sub> (6.37 cm), highest number of leaves (5.50) was observed in T<sub>6</sub> at 30 days after planting and lowest number of leaves (4.40) from treatment T<sub>1</sub> (Control). Shoot length was significant increase in T<sub>6</sub> (6.63 cm) and lowest from T<sub>1</sub> (5.28 cm) at 30 days after planting was recorded. Strawberry plants inoculated with *Azotobacter* + PSB + *Trichoderma* promotes the plant growth due to supplying the essential plant nutrients by the roots. Comparable results were observed in strawberry (Tripathi *et al.*, 2015; Rana and Chandel, 2003; Mishra and Tripathi, 2011; Kumar and Tripathi, 2020). Furthermore, pooled data (Table 2) revealed that,

**Table 1: Effect of plant growth promoting micro-organisms on the plant height, number of leaves and shoot length of strawberry**

Treatments	Plant height (cm)			Number of leaves plant <sup>-1</sup>			Shoot length (cm)		
	30 DAP			30 DAP			30 DAP		
	200-21	2021-22	Pooled	2020-21	2021-22	Pooled	2020-21	2021-22	Pooled
T <sub>1</sub>	5.19	5.12	5.15	4.17	4.63	4.40	5.29	5.26	5.28
T <sub>2</sub>	5.76	5.23	5.49	4.80	4.47	4.63	5.75	5.55	5.65
T <sub>3</sub>	5.54	5.48	5.51	4.73	4.13	4.43	5.43	5.37	5.40
T <sub>4</sub>	5.88	5.77	5.82	4.90	4.73	4.82	5.91	5.88	5.89
T <sub>5</sub>	6.01	5.97	5.99	5.03	4.87	4.95	6.17	6.12	6.14
T <sub>6</sub>	6.15	6.60	6.37	5.20	5.80	5.50	6.35	6.90	6.63
T <sub>7</sub>	6.10	6.40	6.25	5.10	5.13	5.12	6.26	6.19	6.23
T <sub>8</sub>	6.26	7.02	6.64	5.50	5.40	5.45	6.49	6.49	6.49
<b>SEm(±)</b>	<b>0.16</b>	<b>0.08</b>	<b>0.20</b>	<b>0.213</b>	<b>0.31</b>	<b>0.20</b>	<b>0.07</b>	<b>0.14</b>	<b>0.11</b>
<b>LSD(0.05)</b>	<b>0.51</b>	<b>0.25</b>	<b>0.68</b>	<b>0.653</b>	<b>0.94</b>	<b>0.67</b>	<b>0.21</b>	<b>0.43</b>	<b>0.38</b>

**Table 2: Effect of plant growth prompting micro-organisms on the shoot fresh weight, root fresh weight and shoot dry weight of strawberry**

Treatments	Shoot fresh weight (g)			Root fresh weight (g)			Shoot dry weight (g)		
	30 DAP			30 DAP			30 DAP		
	2020-21	2021-22	Pooled	2020-21	2021-22	Pooled	2020-21	2021-22	Pooled
T <sub>1</sub>	25.60	26.21	25.91	7.08	8.26	7.67	5.05	5.21	5.13
T <sub>2</sub>	26.26	26.66	26.46	7.51	8.62	8.07	5.38	5.44	5.41
T <sub>3</sub>	25.87	26.57	26.22	7.34	8.47	7.91	5.21	5.26	5.24
T <sub>4</sub>	26.33	26.96	26.65	7.63	8.73	8.18	5.29	5.50	5.40
T <sub>5</sub>	26.45	27.45	26.95	8.07	8.94	8.51	5.67	5.78	5.72
T <sub>6</sub>	27.94	28.04	27.99	8.89	9.69	9.29	6.21	6.28	6.24
T <sub>7</sub>	26.89	27.59	27.24	8.65	9.52	9.08	5.86	5.94	5.90
T <sub>8</sub>	27.22	27.80	27.51	9.37	9.78	9.57	5.91	6.15	6.03
<b>SEm(±)</b>	<b>0.40</b>	<b>0.32</b>	<b>0.13</b>	<b>0.15</b>	<b>0.14</b>	<b>0.12</b>	<b>0.13</b>	<b>0.13</b>	<b>0.04</b>
<b>LSD(0.05)</b>	<b>1.21</b>	<b>0.99</b>	<b>0.44</b>	<b>0.46</b>	<b>0.42</b>	<b>0.43</b>	<b>0.40</b>	<b>0.41</b>	<b>0.12</b>

**Table 3: Effect of plant growth promoting micro-organisms on root dry weight, number of crowns plant<sup>-1</sup> and estimated fruit yield (t ha<sup>-1</sup>) yield plant<sup>-1</sup> of strawberry**

Treatments	Root dry weight (g)			Number of crowns plant <sup>-1</sup>			Estimated fruit yield (t ha <sup>-1</sup> )		
	2020-21	2021-22	Pooled	2020-21	2021-22	Pooled	2020-21	2021-22	Pooled
T <sub>1</sub>	1.08	1.19	1.14	7.37	7.53	7.45	20.67	21.88	21.27
T <sub>2</sub>	1.81	1.87	1.84	7.73	7.97	7.85	23.06	23.98	23.52
T <sub>3</sub>	1.64	1.76	1.70	7.50	7.60	7.55	21.81	23.38	22.60
T <sub>4</sub>	2.06	2.17	2.12	7.80	8.57	8.18	23.87	24.42	24.15
T <sub>5</sub>	2.26	2.34	2.30	8.17	8.87	8.52	24.74	25.07	24.90
T <sub>6</sub>	2.49	2.72	2.61	8.30	9.63	8.97	26.30	26.69	26.50
T <sub>7</sub>	2.33	2.48	2.40	8.23	9.20	8.72	25.62	25.77	25.70
T <sub>8</sub>	2.67	2.88	2.78	8.43	9.87	9.15	27.06	27.51	27.29
<b>SEm(±)</b>	<b>0.06</b>	<b>0.12</b>	<b>0.03</b>	<b>0.10</b>	<b>0.20</b>	<b>0.26</b>	<b>5.03</b>	<b>6.66</b>	<b>2.44</b>
<b>LSD(0.05)</b>	<b>0.17</b>	<b>0.36</b>	<b>0.10</b>	<b>0.29</b>	<b>0.61</b>	<b>0.88</b>	<b>15.41</b>	<b>20.40</b>	<b>8.31</b>

maximum shoot fresh weight after harvest (27.99 g) was found significantly in T<sub>6</sub> followed by T<sub>8</sub> and T<sub>7</sub> (27.51 g, 27.24 g) respectively. The maximum root fresh weight (9.57 g) was observed from T<sub>6</sub> followed by T<sub>8</sub> (9.57 g). Under T<sub>6</sub> the maximum shoot dry weight (6.24 g) was recorded (Table 2). The maximum root dry weight (2.78 g) in T<sub>8</sub> followed by T<sub>6</sub> (2.61 g) and minimum in T<sub>1</sub> (1.14 g). The treatment T<sub>8</sub> (9.15) recorded maximum crowns per plant than treatment T<sub>1</sub> (7.45). This expansion in vegetative development in plants might be due to the root framework of plant as suggested in strawberry (Tripathi *et al.*, 2010; Kumar and Tripathi, 2020; Singh and Kaur, 2020; Mishra and Tripathi, 2011; Negi *et al.*, 2021).

#### Fruit yield

Pooled data (Table 3) clearly indicated that, estimated fruit yield (27.29 t ha<sup>-1</sup>) was observed maximum with T<sub>8</sub>, followed by T<sub>6</sub> treated plants (26.50 t ha<sup>-1</sup>), while least estimated fruit yield (21.27 t ha<sup>-1</sup>) was recorded under control T<sub>1</sub>. The prolonged photosynthetic capacity of strawberry plants supplied with *Azotobacter* + PSB + *Trichoderma* may have resulted in accumulation of dry matter. Similar findings were obtained in strawberry (Wange *et al.*, 1998; Anonymous, 2022; Tripathi *et al.*, 2010). Biofertilizer containing a living micro-organism that enrich the nutrient status of soil and significantly improves the crop yield.

#### CONCLUSION

Based on the results obtained in the present experiment, performance of strawberry plants exhibits best when treated with *Azotobacter* + PSB + *Trichoderma* (T<sub>8</sub>) as compared to control in order to stimulate growth and yield of strawberry.

#### ACKNOWLEDGEMENT

We acknowledge the support of Uttar Banga Krishi Viswavidyalaya, Pundibari, West Bengal for providing all the facilities to complete the research.

#### REFERENCES

- Anonymous. 2022. National Horticulture Board. Accessed from [nhb.gov.in/report/strawberry](http://nhb.gov.in/report/strawberry)
- Balakrishna, H. T., Swamy, G. S. K., Sabarad, A. I., Patil, P. B., Patil C. P. and Athani, S. I. 2005. Response of *Glomus fasciculatum*, Vermiculture and *Trichoderma harzianum* for yield and yield attributes on banana ratoon crop, *Karnataka J. Hort.*, **1** (4): 63-70.
- Bhagat, P. and Panigrahi, H. 2020. Effect of bio-fertilizer on growth, yield and quality of strawberry (*Fragaria x ananassa* Duch) cv. Nabila under net tunnel, *Pharma Innov. J.*, **9**(1): 442-446.
- Cvijanovi, G., Milosevi, N. and Jarak, M. 2007. The importance of diazotrophs as biofertilizers in the maize and soybean production. *Genetica*, **39** (3): 395-404.
- Esitken, A., Ercisli, S., Karlidag, H. and Sahin, F. 2005. Potential use of Plant Growth Promoting Rhizobacteria (PGPR) in Organic Apricot Production, In: Libek A, Kaufmane E, Sasnauskas A (Eds.), Proceedings of the International Scientific Conference of Environmentally Friendly Fruit Growing. p. 90-97.
- Fernandes, V.C., Domingues, V.F., Mateus, N. and Delerue-Matos, C. 2012. Pesticide residues in Portuguese strawberries grown in 2009-2010 using integrated pest management and organic farming. *Environ. Sci. Pollu. Res.*, **19**:4184-4192.
- Giovannoni, J.J. 2004. Genetic regulation of fruit development and ripening, *Plant Cell*, **16**: 170-180.

- Hazarika, T. K., Nautiyal, B. P. and Bhattacharyya, R. K. 2014. Economic analysis of tissue cultured banana (*Musa paradisiaca* L.) production under the influence of integrated nutrient management, *Indian J. Agric. Sci.*, **84**:656-660.
- Islam, M. T., Hashidoko, Y., Deora, A., Ito, T. and Tahara, S. 2005. Suppression of damping-off disease in host plants by the rhizoplane bacterium *Lysobacter* sp. strain SB-K88 is linked to plant colonization and antibiosis against soil borne Peronosporomycetes. *Appl. Environ. Microbio.*, **71**(7):3786-3796.
- Kumar, A. and Tripathi, V.K. 2020. Effect of *Azotobacter*, PSB and vermicompost on growth, flowering, yield and quality of strawberry (*Fragaria x ananassa* Duch.) cv. Chandler”, *Prog. Hortic.*, **52**(2): 157-161.
- Kumari, S., Chowdhuri, T. K. and Mandal, T. 2018. Effect of biofertilizers on growth and flowering of *Dendrobium var. Sonia*. *J. Crop Weed*, **14**(2): 85-88.
- Mia, M.A.B., Shamsuddin, Z.H., Zakaria, W. and Mahmood, M. 2005. High yielding and quality banana production through plant growth promoting rhizobacterial (PGPR) inoculation, *Fruits Paris*, **60** (3):179-185.
- Mishra, A.N. and Tripathi, V.K. 2011. Influence of different levels of *Azotobacter*, PSB alone and in combination on vegetative growth, flowering, yield and quality of strawberry cv. Chandler, *Int. J. Appl. Agric. Res.*, **6**(3): 203-210.
- Negi, Y.K., Sajwan, P., Uniyal, S. and Mishra, A.C. 2003. Enhancement in yield and nutritive qualities of strawberry fruits by the application of organic manures and biofertilizers, *Scientia Hortic.*, **283**: 1100-1138.
- Rana, R.K. and Chandel, J.S. 2003. Effect of bio-fertilizer and nitrogen on growth, yield and fruit quality of strawberry, *Prog. Hortic.*, **35**(1): 23-30.
- Ruiz, A. F., Portales, B. R., Blanco, M. J. and Caballero, J.L. 2011. The strawberry plant defence mechanism: A molecular review, *Plant Cell Physiol.*, **52**(11): 1873-1903.
- Sabarad, A. I., Swamy, G. S. K., Patil, C. P., Patil, P. B. and Athani, S.I. 2004. Influence of VAM, vermicompost and *Trichoderma harzianum* on growth of banana, cv. Rajapuri (*Musa AAB*)”, *Karnataka J. Agric. Sci.*, **17** (3): 515-518.
- Singh, A. and Singh, J. N. 2009. Effect of bio-fertilizers and bio-regulators on growth, yield and nutrient status of strawberry cv. Sweet Charlie, *Indian J. Hortic.*, **66** (2): 220-224.
- Singh, A.V., Prasad, B. and Shah, S. 2011. Influence of phosphate solubilizing bacteria for enhancement of plant growth and seed yield in lentil. *J. Crop Weed*, **7**(1): 1-4.
- Singh, S. and Kaur, A. 2020. Impact of organic fertilizers on performance of strawberry cv. Sweet Charlie under sub-tropical conditions of Punjab”, *J. Pharmacog. Phytochem.*, **4**: 111-115.
- Tripathi, V.K., Kumar, N., Shukla, H.S. and Mishra, A. N. 2010. Influence of *Azotobacter*, *Azospirillum* and PSB on growth, yield and quality of strawberry cv. Chandler, Paper presented in National Symposium on Conservation Horticulture, during March, 21-23, 2010 at Dehradun, p. 198-199.
- Tripathi, V.K., Kumar, S. and Gupta, A.K. 2015. Influence of *Azotobacter* and vermicompost on growth, flowering, yield and quality of strawberry cv. Chandler, *Indian J. Hortic.*, **72**(2): 201-205.
- Umar, I., Vinod, K.W., Ravi, K. and Mahital, J. 2009. Effect of FYM, urea and *Azotobacter* on growth, yield and quality of strawberry cv. Chandler, *Not. Bot. Hort. Agrobot. Cluj.*, **37** (1): 139-143.
- Wange, S. S., Patil, M. T. and Singh, B. R. 1998. Cultivar × bio-fertilizer interaction study in strawberry, *Recent Hortic.*, **4**: 43-49.