

Studies on chemical weed control in aerobic rice (*Oryza sativa* L.)

P. THIMME GOWDA, *C. SHANKARAIAH, A. C. JNANESH,
M. GOVINDAPPA AND *K. N. MURTHY,

AICRP on Weed Control, MRS, Hebbal, Bangalore

*Department of crop production

Agriculture College, V.C. Farm, Mandy, Karnataka

Aerobic rice production system is gaining importance for increased productivity and reduced water usage and is expected to occupy 10-15 per cent of the total area in India. The major constraints to get higher yield in aerobic rice is weed infestation which cause around 80-90 per cent reduction in grain yield. Manual removal of weeds is supposed to be easy and ecofriendly but highly labour intensive, tedious, back breaking and does not ensure weed removal at critical stages due to non-availability of labours. Hence, there is a need to develop alternative practices for controlling the associated weeds. In such conditions herbicides offer most practical and cost effective means of reducing weed competitions. Therefore, to study the efficacy of some pre-emergence herbicides on aerobic rice the present investigation was undertaken.

The experiment was conducted during *kharif* season of 2005, at Agriculture College, V.C. Farm, Mandy. The soil was sandy loam in texture and slightly acidic in reaction (6.76) with low available nitrogen, medium available phosphorus and potassium. The organic carbon content was medium (0.39 %). Rasi (IET-1444) a popular medium duration variety was sown in mid August with a spacing of 25 x 25 cm. There were included twelve treatments consisted of three doses, each of butachlor (0.75, 1.00 and 1.25 Kg a. i. ha^{-1}), pyrazosulfuron ethyl (20, 25 and 30 g a. i. ha^{-1}) and clomozone + 2,4-DEE (0.75, 1.00 and 1.25 litre ha^{-1}), two hand weeding at 20 and 45 DAS, two inter cultivation at 20 and 45 DAS and weedy check were laid out in Randomized Complete Block Design (RCBD) with three replications. Pre-emergence application of herbicides was done at one day after sowing. Irrespective of the treatments one intercultural operation was given at 45 DAS. The data was subjected to square root transformation using the formula $\sqrt{x}+0.5$ and the statistical analysis was done.

The major weed flora observed in experimental plots were: *Digitaria sanguinalis*, *Cynodon dactylon*, *Panicum repens* and *Dactyloctenium aegyptium*. The narrow leaved weeds (NLW) were, *Aegeratum conyzoids*, *Commelina benghalensis*, *Euphorbia hirta* and *Tridax procumbens*. *Phyllanthus niruri* and *Celosia argentea* were broad leaved weeds (BLW) and *Cyperus rotundus* and *Cyperus iria* were sedges.

Hand weeding at 20 and 45 DAS excelled with a lowest weed population of weeds ($43.67 m^{-2}$) and dry weight of weeds ($3.42 g 0.25 m^{-2}$) among the various treatments. It was at par with clomozone + 2,4-DEE @ 1.25 litre ha^{-1} and pyrazosulfuron ethyl @ 30 g a. i. ha^{-1} for controlling NLW and with pyrazosulfuron ethyl @ 30 g a. i. ha^{-1} for controlling BLW. Among the various herbicides, lowest weed population ($48 m^{-2}$) and dry weight of weeds ($3.94 g 0.25 m^{-2}$) were observed with pyrazosulfuron ethyl @ 30 g a. i. ha^{-1} followed by clomozone + 2,4-DEE @ 1.25 litre ha^{-1} . The highest weed population ($366.33 m^{-2}$) and dry weight of weeds ($43.11 g 0.25 m^{-2}$) were observed with unweeded check. The weed control efficiency was higher with hand weeding (92.07 %) and pyrazosulfuron ethyl @ 30 g a. i. ha^{-1} (90.86 %).

Hand weeding at 20 and 45 DAS recorded significantly taller plant height and higher dry matter production ($72.53 cm$ and $66.25 g hill^{-1}$, respectively) among the various treatments. It was statistically on par with herbicide treatment pyrazosulfuron ethyl @ 30 g a. i. ha^{-1} ($71.53 cm$ and $65.37 g hill^{-1}$). Sharma *et al.* 2004 and Moorthy 2002 also reported similar results in direct seeded rice. Unweeded check registered significantly least plant height and dry matter production ($52.33 cm$ and $16.58 g hill^{-1}$) as a consequence of severe competition of rice plant with weeds for available resources.

The yield attributing parameters *viz*, number of effective tillers per hill, panicle length, filled grains per panicle and 1000 grain weight were found significantly higher with two hand weeding at 20 and 45 DAS ($20, 23.67 cm, 132.73$ and $24.93 g$, respectively) and was remaining statistically at par with pyrazosulfuron ethyl @ 30 g a. i. ha^{-1} . The increase in yield attributing parameters in above mentioned treatments was mainly due to better crop growth. In general higher doses of various herbicides expressed higher growth and yield attributing characters as compared to their lower doses due to their higher weed control efficiency except clomozone + 2,4-DEE (Table 2).

All the weed control treatments registered significantly higher yield than weedy check. The highest grain and straw yield among the treatments was recorded under hand weeding at 20 and 45 DAS

(5.07 t ha⁻¹ and 5.53 t ha⁻¹). It was on par with pyrazosulfuron ethyl @ 30 g a.i. ha⁻¹. The results are in good agreement with like, Budhar *et al.* (1991), Moorthy (1997b) and Moorthy (2002). Unwedded check recorded 82.84 and 82.60 per cent reduction in grain yield and 75.91 and 75.69 per cent reduction in straw yield as compared to hand weeding at 20 and 45 DAS and pyrazosulfuron ethyl @ 30 g a.i. ha⁻¹, respectively. This was due to less number of effective tillers per hill, panicle length, filled grains per panicle and 1000 grain weight. Greater competitions offered by weeds throughout crop growth period suppressed the crop, severely affecting plant height and dry matter production per hill led to the poor yield components and thus lower grain yield.

Hand weeding required additional investment of Rs. 2500 ha⁻¹. for removing weeds. All the herbicidal treatments need lesser additional investment (Rs. 625 to 1062 ha⁻¹) depending upon the cost and rate of herbicide application. The maximum gross returns was (Rs. 31,203 ha⁻¹) hand weeding which was closely followed by of pyrazosulfuron ethyl @ 30 g a.i. ha⁻¹ (Rs. 30,782 ha⁻¹). The net returns and benefit cost ratio was maximum with pyrazosulfuron ethyl @ 30 g a.i. ha⁻¹ (Rs. 21,019 ha⁻¹ and 2.15). The net returns and benefit cost ratio were quite lower (Rs. 10,940 ha⁻¹ and 1.77) under two hand weeding at 20 and 45 DAS which indicated that it was less remunerative than most of the herbicidal weed control treatments, confirming the view of Singh and Govindra Singh (2001).

The above study concludes that hand weeding at 20 and 45 DAS found effective in control of weeds and recorded lower weed population and dry weight among various treatments. It was on par with herbicide treatment pyrazosulfuron ethyl @ 30 g a.i. ha⁻¹. The growth and yield attributing characters were recorded higher with hand weeding at 20 and 45 DAS however, on par with pyrazosulfuron ethyl @ 30 g a.i. ha⁻¹. Hand weeding at 20 and 45 DAS recorded significantly higher grain yield and was at par with herbicide treatment pyrazosulfuron ethyl @ 30 g a.i. ha⁻¹.

REFERENCES

Budhar, M.N., Munoli, S. and Ram S. C., 1991, Weed control in direct sown rice under puddled condition. *Indian J. Weed Sci.*, **23** ; 76-72.

Moorthy, B.T.S., 1997b, Chemical weed control in puddle seeded rice, *Oryza*, **29**: 54-58.

Moorthy, B.T.S., 2002, Evaluation of pyrazosulfuron ethyl alone and in combination with molinate for controlling weeds in rainfed direct seeded low land rice. *Indian J. Weed Sci.*, **34**: 79-80.

Sharma, S.D., Punia, S.S. Narwal, S. And Malik, P.K., 2004, Effects of different rate and time of application of Pyrazosulfuron ethyl against weeds in transplanted rice. *Haryana J. Agron.*, **20** : 23-25.

Singh, V.P. And Singh, G. 2001, Weed control studies in spring rice (*Oryza sativa* L.) under rainfed low valley situation of Uttaranchal. *Indian J. Weed Sci.*, **33**: 52-55.

Table 1: Effect of weed control treatments on weed population, dry weight and weed control efficiency (WCE)

Treatments	Weed population (No. m ⁻²) at harvest								Weed dry weight (g 0.25 m ⁻²) at harvest				WCE (%)		
	NLW		BLW		Sedges		Total weeds		Grasses		BLW		Sedges		
	NLW	BLW	Sedges	Total weeds	Grasses	BLW	Sedges	Total weeds							
T ₁	5.15 (26.33)	9.61 (92.00)	4.92 (23.67)	11.93 (142.00)	2.91 (8.00)	3.23 (10.00)	2.16 (4.16)	4.82 (22.16)	48.60						
T ₂	4.49 (20.00)	7.36 (54.33)	3.78 (14.00)	9.30 (88.33)	2.35 (5.00)	2.80 (7.40)	1.64 (2.10)	4.00 (14.59)	66.16						
T ₃	3.98 (12.67)	6.75 (45.00)	3.66 (13.00)	8.39 (70.67)	2.12 (4.00)	2.47 (5.83)	1.50 (1.75)	3.54 (11.58)	73.14						
T ₄	5.58 (31.00)	7.70 (59.00)	1.92 (3.33)	9.66 (93.33)	3.21 (9.80)	2.30 (4.80)	0.83 (0.19)	3.96 (14.79)	65.69						
T ₅	4.40 (19.00)	6.99 (48.67)	0.89 (0.33)	8.27 (68.00)	2.71 (6.83)	1.90 (3.10)	0.71 (0.00)	3.23 (9.93)	76.97						
T ₆	3.47 (11.67)	6.06 (36.33)	0.71 (0.00)	6.96 (48.00)	1.43 (1.54)	1.70 (2.40)	0.71 (0.00)	2.10 (3.94)	90.86						
T ₇	4.33 (19.33)	7.47 (55.67)	5.00 (24.67)	9.96 (99.67)	1.74 (2.67)	3.05 (8.83)	2.55 (6.00)	4.30 (17.50)	59.41						
T ₈	3.61 (13.33)	6.56 (43.00)	3.84 (14.33)	8.39 (70.66)	1.44 (1.63)	2.55 (6.00)	2.11 (3.95)	3.57 (11.58)	73.14						
T ₉	3.05 (9.33)	6.51 (42.33)	3.78 (13.67)	8.08 (65.33)	1.21 (1.00)	2.40 (5.33)	1.89 (3.07)	3.25 (9.40)	78.20						
T ₁₀	4.48 (19.67)	7.35 (54.00)	3.87 (14.67)	9.40 (88.34)	2.53 (5.93)	2.86 (7.73)	1.89 (3.07)	4.18 (16.73)	61.19						
T ₁₁	2.11 (4.00)	5.92 (34.67)	2.32 (5.00)	6.66 (43.67)	1.16 (0.87)	1.51 (1.80)	1.12 (0.75)	1.86 (3.42)	92.07						
T ₁₂	7.27 (53.33)	15.71 (247.33)	8.13 (65.67)	19.19 (366.33)	3.97 (15.33)	4.32 (18.17)	3.18 (9.61)	6.67 (43.11)	-						
S.Em. \pm	0.454	0.389	0.190	0.507	0.127	0.161	0.113	0.177	-						
LSD (P=0.05)	1.331	1.141	0.557	1.484	0.371	0.472	0.331	0.519							

Values in the parenthesis are original values

T₁: Butachlor @ 0.75 kg a.i. ha⁻¹T₃: Butachlor @ 1.25 kg a.i. ha⁻¹T₅: Pyrazosulfuron ethyl @ 25 g a.i. ha⁻¹T₇: Clomozone + 2,4 - DEE (RM) @ 0.75 lit. ha⁻¹T₉: Clomozone + 2,4 - DEE (RM) @ 1.25 lit. ha⁻¹T₁₁: Two HW at 20 & 45 DAST₂: Butachlor @ 1.00 kg a.i. ha⁻¹T₄: Pyrazosulfuron ethyl @ 20 g a.i. ha⁻¹T₆: Pyrazosulfuron ethyl @ 30 g a.i. ha⁻¹T₈: Clomozone + 2,4 - DEE (RM) @ 1.00 lit. ha⁻¹T₁₀: Two IC at 20 & 45 DAST₁₂: Weedy check

BLW: Broad leaved weeds

NLW :Narrow leaved weeds

RM : Ready Mix

IC : Inter Cultivation

HW : Hand Weeding

Table 2: Effect of weed control treatments on growth, yield and yield parameters of aerobic rice

Treatments	Plant height (cm)	Dry matter (g hill ⁻¹)	No. of effective tillers hill ⁻¹	Panicle length (cm)	Filled grains per panicle	1000 grain weight (g)	Grain yield (t ha ⁻¹)	Straw yield (t ha ⁻¹)
T ₁	56.97	37.40	10.33	18.57	95.47	22.33	2.76	3.71
T ₂	62.90	48.01	15.60	20.50	111.40	23.17	3.37	4.16
T ₃	64.27	52.50	16.80	20.90	116.53	23.83	3.75	4.49
T ₄	59.20	40.89	13.33	18.60	98.20	23.17	2.95	3.86
T ₅	65.43	52.63	15.47	20.00	110.87	23.97	3.81	4.61
T ₆	71.53	65.37	19.30	23.30	132.27	24.83	5.00	5.47
T ₇	62.33	42.19	13.13	19.17	104.73	23.23	2.96	3.79
T ₈	67.10	56.62	17.40	21.00	117.80	24.17	4.16	4.90
T ₉	66.93	52.31	16.57	20.80	113.80	24.00	3.81	4.56
T ₁₀	60.50	45.79	15.13	19.57	113.07	22.33	3.23	4.07
T ₁₁	72.53	66.25	20.00	23.67	132.73	24.93	5.07	5.53
T ₁₂	52.33	16.58	4.67	16.33	54.47	21.30	0.87	1.33
S.Em. \pm	1.477	1.517	0.573	0.703	3.953	0.441	0.167	0.163
C.D.= P=0.05) at 5%	4.332	4.451	1.680	2.062	11.594	1.299	0.489	0.478

T₁: Butachlor @ 0.75 kg a.i. ha⁻¹T₂: Butachlor @ 1.00 kg a.i. ha⁻¹

BLW: Broad leaved weeds

T₃: Butachlor @ 1.25 kg a.i. ha⁻¹T₄: Pyrazosulfuron ethyl @ 20 g a.i. ha⁻¹

NLW :Narrow leaved weeds

T₅: Pyrazosulfuron ethyl @ 25 g a.i. ha⁻¹T₆: Pyrazosulfuron ethyl @ 30 g a.i. ha⁻¹

RM : Ready Mix

T₇: Clomozone + 2,4 - DEE (RM) @ 0.75 lit. ha⁻¹T₈: Clomozone + 2,4 - DEE (RM) @ 1.00 lit. ha⁻¹

IC : Inter Cultivation

T₉: Clomozone + 2,4 - DEE (RM) @ 1.25 lit. ha⁻¹T₁₀: Two IC at 20 & 45 DAS

HW : Hand Weeding

T₁₁: Two HW at 20 & 45 DAST₁₂: Weedy check**Table 3: Economics of weed control treatments in aerobic rice**

Treatments	Cost of weed control (Rs. ha ⁻¹)	Total cost of cultivation (Rs. ha ⁻¹)	Gross returns (Rs. ha ⁻¹)	Net returns (Rs. ha ⁻¹)	Benefit: Cost ratio
	625	9388	17406	8018	0.85
	750	9513	21031	11518	1.21
T ₃	875	9638	23319	13681	1.42
T ₄	750	9513	18541	9028	0.95
T ₅	875	9638	23721	14083	1.46
T ₆	1000	9763	30782	21019	2.15
T ₇	737	9500	18554	9054	0.95
T ₈	900	9663	25820	18157	1.88
T ₉	1062	9825	23691	13866	1.41
T ₁₀	600	9363	20207	10844	1.16
T ₁₁	2500	11263	31203	10940	1.77
T ₁₂	-	8763	5583	-3180	-0.36